arXiv:1407.3550v2 [math.NT] 18 Aug 2014

Dedekind η-function, Hauptmodul and invariant theory

Lei Yang

Abstract

We solve a long-standing open problem with its own long history dating back to the celebrated works of Klein and Ramanujan. This problem concerns the invariant decomposition formulas of the Hauptmodul for $\Gamma_0(p)$ under the action of finite simple groups $PSL(2,p)$ with $p = 5, 7, 13$. The cases of $p = 5$ and 7 were solved by Klein and Ramanujan. Little was known about this problem for $p = 13$. Using our invariant theory for $PSL(2, 13)$, we solve this problem. This leads to a new expression of the classical elliptic modular function of Klein: j-function in terms of theta constants associated with $\Gamma(13)$. Moreover, we find an exotic modular equation, i.e., it has the same form as Ramanujan's modular equation of degree 13, but with different kinds of modular parametrizations, which gives the geometry of the classical modular curve $X(13)$.

Contents

1. Introduction
2. Six-dimensional representations of $PSL(2, 13)$ and transformation formulas for theta constants
3. Seven-dimensional representations of $PSL(2, 13)$, exotic modular equation and geometry of modular curve $X(13)$
4. Fourteen-dimensional representations of $PSL(2, 13)$ and invariant decomposition formula

1. Introduction

In many applications of elliptic modular functions to number theory the Dedekind eta function plays a central role. It is defined in the upper-half plane $\mathbb{H} = \{z \in \mathbb{C} : \mathrm{Im}(z) > 0\}$ by

$$\eta(z) := q^{\frac{1}{24}} \prod_{n=1}^{\infty}(1 - q^n), \quad q = e^{2\pi iz}.$$

Typeset by $\mathcal{AMS}$-TEX

The Dedekind eta function is closely related to the partition function. A partition of a positive integer n is any non-increasing sequence of positive integers whose sum is n. Let $p(n)$ denote the number of partitions of n. The partition function $p(n)$ has the well-known generating function

$$\sum_{n=0}^{\infty} p(n)q^n = \prod_{n=1}^{\infty} \frac{1}{1-q^n} = \frac{q^{\frac{1}{24}}}{\eta(z)}.$$

In his ground-breaking works [R1], [R2] and [R3], Ramanujan found the following famous Ramanujan partition congruences:

$$\begin{aligned} p(5n+4) &\equiv 0 \pmod 5, \\ p(7n+5) &\equiv 0 \pmod 7, \\ p(11n+6) &\equiv 0 \pmod{11}. \end{aligned}$$

Ramanujan's proofs of the congruences modulo 5 and 7 are quite ingenious but are not terribly difficult:

$$\begin{aligned} q^{\frac{19}{24}} \sum_{n=0}^{\infty} p(5n+4)q^n &= 5\frac{\eta(5z)^5}{\eta(z)^6}, \\ q^{\frac{17}{24}} \sum_{n=0}^{\infty} p(7n+5)q^n &= 7\frac{\eta(7z)^3}{\eta(z)^4} + 49\frac{\eta(7z)^7}{\eta(z)^8}, \end{aligned} \tag{1.1}$$

while the proof of the congruence modulo 11 is much harder. One of the significance about Ramanujan partition congruences comes from the problem of classifying congruences of the form $p(ln+b) \equiv 0 \pmod l$, where l is a prime. It was proved in [AB] that there are only those three originally observed by Ramanujan, when $(l, b) = (5, 4)$, $(7, 5)$, or $(11, 6)$.

In order to provide a combinatorial explanation of these congruences, Dyson in [D] defined the rank of a partition to be its largest part minus the number of its parts. He conjectured that the partitions of $5n + 4$ (resp. $7n + 5$) are divided into 5 (resp. 7) groups of equal size when sorted by their ranks modulo 5 (resp. 7), thereby providing a combinatorial explanation for the congruences mod 5 and 7. This conjecture was proved by Atkin and Swinnerton-Dyer (see [ASD]). However, for the third congruence, the corresponding criterion failed, and so Dyson conjectured the existence of a statistic, which he called the crank to combinatorially explain the congruence $p(11n+6) \equiv 0 \pmod{11}$. The crank of a partition was found by Andrews and Garvan (see [AG]). The crank divides the partition into equinumerous congruence classes modulo 5, 7 and 11 for the three congruences, respectively. Further work of Garvan, Kim and Stanton (see [GKS]) produces, for the congruences with moduli 5, 7, 11 and 25, combinatorial interpretations which are rooted in the modular representation theory of the symmetric group. In fact,

in his lost notebook [R], Ramanujan had recorded the generating functions for both the rank and the crank. Moreover, he actually recorded an observation that is equivalent to Dyson's assertion about the congruences for the ranks of the partitions of $5n+4$ (see [G1] and [BCCL]). Recently, Bringmann and Ono (see [BrO]) showed that the rank generating functions are related to harmonic weak Maass forms. Using Shimura's theory of half-integer weight modular forms and Serre's theory of p-adic modular forms, they obtained as an application infinite families of congruences for the rank. These generalize previous work of Ono (see [O]) on the partition function. In [Ma], Mahlburg developed an analogous theory of congruences for crank generating functions, which are shown to possess the same sort of arithmetic properties.

In [Z] Zuckerman found an identity in the spirit of (1.1) (see also [R] and [BO]):

$$\begin{aligned} q^{\frac{11}{24}}\sum_{n=0}^{\infty}p(13n+6)q^n =& 11\frac{\eta(13z)}{\eta(z)^2}+36\cdot 13\frac{\eta(13z)^3}{\eta(z)^4}+38\cdot 13^2\frac{\eta(13z)^5}{\eta(z)^6}+20\cdot 13^3\frac{\eta(13z)^7}{\eta(z)^8} \\ &+6\cdot 13^4\frac{\eta(13z)^9}{\eta(z)^{10}}+13^5\frac{\eta(13z)^{11}}{\eta(z)^{12}}+13^5\frac{\eta(13z)^{13}}{\eta(z)^{14}}. \end{aligned} \tag{1.2}$$

Rademacher (see [Ra]) pointed out that (1.1) can be rewritten as

$$\begin{aligned} \sum_{\lambda=0}^{4}\eta(5z)\eta\left(\frac{z+24\lambda}{5}\right)^{-1} &= 5^2\left(\frac{\eta(5z)}{\eta(z)}\right)^6, \\ \sum_{\lambda=0}^{6}\eta(7z)\eta\left(\frac{z+24\lambda}{7}\right)^{-1} &= 7^2\left(\frac{\eta(7z)}{\eta(z)}\right)^4+7^3\left(\frac{\eta(7z)}{\eta(z)}\right)^8. \end{aligned} \tag{1.3}$$

In fact, (1.2) can also be rewritten as

$$\begin{aligned} \sum_{\lambda=0}^{12}\eta(13z)\eta\left(\frac{z+24\lambda}{13}\right)^{-1} &= 11\cdot 13\left(\frac{\eta(13z)}{\eta(z)}\right)^2+ \\ &+36\cdot 13^2\left(\frac{\eta(13z)}{\eta(z)}\right)^4+38\cdot 13^3\left(\frac{\eta(13z)}{\eta(z)}\right)^6+20\cdot 13^4\left(\frac{\eta(13z)}{\eta(z)}\right)^8+ \\ &+6\cdot 13^5\left(\frac{\eta(13z)}{\eta(z)}\right)^{10}+13^6\left(\frac{\eta(13z)}{\eta(z)}\right)^{12}+13^6\left(\frac{\eta(13z)}{\eta(z)}\right)^{14}. \end{aligned} \tag{1.4}$$

Let $X=X_0(p)$ be the compactification of $\mathbb{H}/\Gamma_0(p)$ where p is prime. The complex function field of X consists of the modular functions $f(z)$ for $\Gamma_0(p)$ which are meromorphic on the extended upper half-plane. A function f lies in the rational function field $\mathbb{Q}(X)$ if and only if the Fourier coefficients in its expansion at ∞: $f(z)=\sum a_n q^n$ are

all rational numbers. The field $\mathbb{Q}(X)$ is known to be generated over $\mathbb{Q}$ by the classical j-functions

$$\begin{cases} j = j(z) = q^{-1} + 744 + 196884q + \cdots, \\ j_p = j\left(\dfrac{-1}{pz}\right) = j(pz) = q^{-p} + 744 + \cdots. \end{cases}$$

A further element in the function field $\mathbb{Q}(X) = \mathbb{Q}(j, j_p)$ is the modular unit $u = \Delta(z)/\Delta(pz)$ with divisor $(p-1)\{(0) - (\infty)\}$ where $\Delta(z)$ is the discriminant. If $m = \gcd(p-1, 12)$, then an mth root of u lies in $\mathbb{Q}(X)$. This function has the Fourier expansion

$$t = \sqrt[m]{u} = q^{(1-p)/m} \prod_{n\geq 1} \left(\frac{1-q^n}{1-q^{np}}\right)^{24/m} = \left(\frac{\eta(z)}{\eta(pz)}\right)^{24/m}.$$

When $p-1$ divides 12, so $m = p-1$, the function t is a Hauptmodul for the curve X which has genus zero (see [Gr]). It is well-known that the genus of the modular curve X for prime p is zero if and only if $p = 2, 3, 5, 7, 13$. In his paper [K1], Klein studied the modular equations of orders 2, 3, 5, 7, 13 with degrees 3, 4, 6, 8, 14, respectively. They are the so-called Hauptmoduln (principal moduli) which means that there is a uniformising function J_Γ that is a modular function for some congruence subgroup Γ, and all other modular functions for Γ can be written as a rational function in it. Hence, (1.3) and (1.4) are intimately related to the Hauptmoduln for modular curves $X_0(5)$, $X_0(7)$ and $X_0(13)$.

The modular curves $X_0(5)$ and $X_0(7)$ were studied by Klein in his pioneering work (see [K], [K1], [K2], [KF1] and [KF2]) and by Ramanujan (see [R5], [R6] and [R]). It is well-known that the celebrated Rogers-Ramanujan identities

$$G(q) := \sum_{n=0}^{\infty} \frac{q^{n^2}}{(1-q)\cdots(1-q^n)} = \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+1})(1-q^{5n+4})}$$

and

$$H(q) := \sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(1-q)\cdots(1-q^n)} = \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+2})(1-q^{5n+3})}$$

are intimately associated with the Rogers-Ramanujan continued fraction

$$R(q) := \cfrac{q^{\frac{1}{5}}}{1 + \cfrac{q}{1 + \cfrac{q^2}{1 + \cfrac{q^3}{1 + \cdots}}}},$$

namely, they satisfy $R(q) = q^{\frac{1}{5}} \frac{H(q)}{G(q)}$. In [R4], Ramanujan found an algebraic relation between $G(q)$ and $H(q)$:

$$G^{11}(q)H(q) - q^2 G(q)H^{11}(q) = 1 + 11qG^6(q)H^6(q),$$

which is equivalent to one of the most important formulas for $R(q)$ (see also [AB1]):

$$\frac{1}{R^5(q)} - 11 - R^5(q) = \left(\frac{\eta(z)}{\eta(5z)}\right)^6. \tag{1.5}$$

In his celebrated work on elliptic modular functions (see [KF1], p.640, [KF2], p.73 and [Du]), Klein showed that $R(q) = \frac{a(z)}{b(z)}$ where

$$a(z) = e^{-\frac{3\pi i}{10}} \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, 5z), \quad b(z) = e^{-\frac{\pi i}{10}} \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, 5z).$$

Hence, (1.5) is equivalent to the following formula

$$\left(\frac{\eta(z)}{\eta(5z)}\right)^6 = -\frac{f(a(z), b(z))}{a(z)^6 b(z)^6}, \tag{1.6}$$

where

$$f(z_1, z_2) = z_1 z_2 (z_1^{10} + 11 z_1^5 z_2^5 - z_2^{10})$$

is an invariant of degree 12 associated to the icosahedron, i.e., it is invariant under the action of the simple group $PSL(2, 5)$, and $z_1^2 z_2^2$ is invariant under the action of the image of a Borel subgroup of $PSL(2, 5)$ (i.e. a maximal subgroup of order 10 of $PSL(2, 5)$) (see [K] for more details). We call (1.6) the invariant decomposition formula for the icosahedron.

In his work on elliptic modular functions (see [KF1], p.746), Klein also obtained the invariant decomposition formula for the simple group $PSL(2, 7)$ of order 168:

$$\left(\frac{\eta(z)}{\eta(7z)}\right)^4 = \frac{\Phi_6(a(z), b(z), c(z))}{a(z)^2 b(z)^2 c(z)^2}, \tag{1.7}$$

where

$$a(z) = -e^{-\frac{5\pi i}{14}} \theta \begin{bmatrix} \frac{5}{7} \\ 1 \end{bmatrix} (0, 7z), \quad b(z) = e^{-\frac{3\pi i}{14}} \theta \begin{bmatrix} \frac{3}{7} \\ 1 \end{bmatrix} (0, 7z), \quad c(z) = e^{-\frac{\pi i}{14}} \theta \begin{bmatrix} \frac{1}{7} \\ 1 \end{bmatrix} (0, 7z),$$

and

$$\Phi_6(x, y, z) = xy^5 + yz^5 + zx^5 - 5x^2 y^2 z^2$$

is an invariant of degree 6 associated to $PSL(2,7)$, and xyz is invariant under the action of the image of a Borel subgroup of $PSL(2,7)$ (i.e. a maximal subgroup of order 21 of $PSL(2,7)$) (see [K2] for more details). Independently, Ramanujan also gave the same formula (see [R6], p.300, [B2], p.174, [El] and [La] for more details).

The invariant decomposition formula for the simple group $PSL(2,13)$ of order 1092 is much harder and the modular curve $X_0(13)$ is much more complicated. In his monograph (see [KF2], p.73), Klein showed that $(\eta(z)/\eta(13z))^2$ is a Hauptmodul for $\Gamma_0(13)$. Ramanujan also studied this problem (see [R5], p.326, [R6], p.244 and [B1], p.372). However, neither of them could obtain the invariant decomposition formula in the spirit of (1.6) and (1.7) (see the end of section 4 for more details).

In the present paper, we establish the invariant theory for $PSL(2,13)$. Combining with theta constants for $\Gamma(13)$, we obtain the invariant decomposition formula for $PSL(2,13)$. Let

$$\begin{cases} a_1(z) := e^{-\frac{11\pi i}{26}}\theta\begin{bmatrix}\frac{11}{13}\\ 1\end{bmatrix}(0,13z) = q^{\frac{121}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+11n)}, \\ a_2(z) := e^{-\frac{7\pi i}{26}}\theta\begin{bmatrix}\frac{7}{13}\\ 1\end{bmatrix}(0,13z) = q^{\frac{49}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+7n)}, \\ a_3(z) := e^{-\frac{5\pi i}{26}}\theta\begin{bmatrix}\frac{5}{13}\\ 1\end{bmatrix}(0,13z) = q^{\frac{25}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+5n)}, \\ a_4(z) := -e^{-\frac{3\pi i}{26}}\theta\begin{bmatrix}\frac{3}{13}\\ 1\end{bmatrix}(0,13z) = -q^{\frac{9}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+3n)}, \\ a_5(z) := e^{-\frac{9\pi i}{26}}\theta\begin{bmatrix}\frac{9}{13}\\ 1\end{bmatrix}(0,13z) = q^{\frac{81}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+9n)}, \\ a_6(z) := e^{-\frac{\pi i}{26}}\theta\begin{bmatrix}\frac{1}{13}\\ 1\end{bmatrix}(0,13z) = q^{\frac{1}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+n)} \end{cases} \tag{1.8}$$

be the theta constants for $\Gamma(13)$. Set

$$\Phi_{12}(z_1,z_2,z_3,z_4,z_5,z_6) := -\frac{1}{26}(7\cdot 13^2\mathbb{G}_0^2 + \mathbb{G}_1\mathbb{G}_{12} + \mathbb{G}_2\mathbb{G}_{11} + \cdots + \mathbb{G}_6\mathbb{G}_7), \tag{1.9}$$

where

$$\begin{cases}
\mathbb{G}_0 = \mathbb{D}_0^2 + \mathbb{D}_\infty^2, \\
\mathbb{G}_1 = -\mathbb{D}_7^2 + 2\mathbb{D}_0\mathbb{D}_1 + 10\mathbb{D}_\infty\mathbb{D}_1 + 2\mathbb{D}_2\mathbb{D}_{12} - 2\mathbb{D}_3\mathbb{D}_{11} - 4\mathbb{D}_4\mathbb{D}_{10} - 2\mathbb{D}_9\mathbb{D}_5, \\
\mathbb{G}_2 = -2\mathbb{D}_1^2 - 4\mathbb{D}_0\mathbb{D}_2 + 6\mathbb{D}_\infty\mathbb{D}_2 - 2\mathbb{D}_4\mathbb{D}_{11} + 2\mathbb{D}_5\mathbb{D}_{10} - 2\mathbb{D}_6\mathbb{D}_9 - 2\mathbb{D}_7\mathbb{D}_8, \\
\mathbb{G}_3 = -\mathbb{D}_8^2 + 2\mathbb{D}_0\mathbb{D}_3 + 10\mathbb{D}_\infty\mathbb{D}_3 + 2\mathbb{D}_6\mathbb{D}_{10} - 2\mathbb{D}_9\mathbb{D}_7 - 4\mathbb{D}_{12}\mathbb{D}_4 - 2\mathbb{D}_1\mathbb{D}_2, \\
\mathbb{G}_4 = -\mathbb{D}_2^2 + 10\mathbb{D}_0\mathbb{D}_4 - 2\mathbb{D}_\infty\mathbb{D}_4 + 2\mathbb{D}_5\mathbb{D}_{12} - 2\mathbb{D}_9\mathbb{D}_8 - 4\mathbb{D}_1\mathbb{D}_3 - 2\mathbb{D}_{10}\mathbb{D}_7, \\
\mathbb{G}_5 = -2\mathbb{D}_9^2 - 4\mathbb{D}_0\mathbb{D}_5 + 6\mathbb{D}_\infty\mathbb{D}_5 - 2\mathbb{D}_{10}\mathbb{D}_8 + 2\mathbb{D}_6\mathbb{D}_{12} - 2\mathbb{D}_2\mathbb{D}_3 - 2\mathbb{D}_{11}\mathbb{D}_7, \\
\mathbb{G}_6 = -2\mathbb{D}_3^2 - 4\mathbb{D}_0\mathbb{D}_6 + 6\mathbb{D}_\infty\mathbb{D}_6 - 2\mathbb{D}_{12}\mathbb{D}_7 + 2\mathbb{D}_2\mathbb{D}_4 - 2\mathbb{D}_5\mathbb{D}_1 - 2\mathbb{D}_8\mathbb{D}_{11}, \\
\mathbb{G}_7 = -2\mathbb{D}_{10}^2 + 6\mathbb{D}_0\mathbb{D}_7 + 4\mathbb{D}_\infty\mathbb{D}_7 - 2\mathbb{D}_1\mathbb{D}_6 - 2\mathbb{D}_2\mathbb{D}_5 - 2\mathbb{D}_8\mathbb{D}_{12} - 2\mathbb{D}_9\mathbb{D}_{11}, \\
\mathbb{G}_8 = -2\mathbb{D}_4^2 + 6\mathbb{D}_0\mathbb{D}_8 + 4\mathbb{D}_\infty\mathbb{D}_8 - 2\mathbb{D}_3\mathbb{D}_5 - 2\mathbb{D}_6\mathbb{D}_2 - 2\mathbb{D}_{11}\mathbb{D}_{10} - 2\mathbb{D}_1\mathbb{D}_7, \\
\mathbb{G}_9 = -\mathbb{D}_{11}^2 + 2\mathbb{D}_0\mathbb{D}_9 + 10\mathbb{D}_\infty\mathbb{D}_9 + 2\mathbb{D}_5\mathbb{D}_4 - 2\mathbb{D}_1\mathbb{D}_8 - 4\mathbb{D}_{10}\mathbb{D}_{12} - 2\mathbb{D}_3\mathbb{D}_6, \\
\mathbb{G}_{10} = -\mathbb{D}_5^2 + 10\mathbb{D}_0\mathbb{D}_{10} - 2\mathbb{D}_\infty\mathbb{D}_{10} + 2\mathbb{D}_6\mathbb{D}_4 - 2\mathbb{D}_3\mathbb{D}_7 - 4\mathbb{D}_9\mathbb{D}_1 - 2\mathbb{D}_{12}\mathbb{D}_{11}, \\
\mathbb{G}_{11} = -2\mathbb{D}_{12}^2 + 6\mathbb{D}_0\mathbb{D}_{11} + 4\mathbb{D}_\infty\mathbb{D}_{11} - 2\mathbb{D}_9\mathbb{D}_2 - 2\mathbb{D}_5\mathbb{D}_6 - 2\mathbb{D}_7\mathbb{D}_4 - 2\mathbb{D}_3\mathbb{D}_8, \\
\mathbb{G}_{12} = -\mathbb{D}_6^2 + 10\mathbb{D}_0\mathbb{D}_{12} - 2\mathbb{D}_\infty\mathbb{D}_{12} + 2\mathbb{D}_2\mathbb{D}_{10} - 2\mathbb{D}_1\mathbb{D}_{11} - 4\mathbb{D}_3\mathbb{D}_9 - 2\mathbb{D}_4\mathbb{D}_8
\end{cases} \tag{1.10}$$

are the senary sextic forms (sextic forms in six variables). Here,

$$\begin{cases}
\mathbb{D}_0 = z_1z_2z_3, \\
\mathbb{D}_1 = 2z_2z_3^2 + z_2^2z_6 - z_4^2z_5 + z_1z_5z_6, \\
\mathbb{D}_2 = -z_6^3 + z_2^2z_4 - 2z_2z_5^2 + z_1z_4z_5 + 3z_3z_5z_6, \\
\mathbb{D}_3 = 2z_1z_2^2 + z_1^2z_5 - z_4z_6^2 + z_3z_4z_5, \\
\mathbb{D}_4 = -z_2^2z_3 + z_1z_6^2 - 2z_4^2z_6 - z_1z_3z_5, \\
\mathbb{D}_5 = -z_4^3 + z_3^2z_5 - 2z_3z_6^2 + z_2z_5z_6 + 3z_1z_4z_6, \\
\mathbb{D}_6 = -z_5^3 + z_1^2z_6 - 2z_1z_4^2 + z_3z_4z_6 + 3z_2z_4z_5, \\
\mathbb{D}_7 = -z_2^3 + z_3z_4^2 - z_1z_3z_6 - 3z_1z_2z_5 + 2z_1^2z_4, \\
\mathbb{D}_8 = -z_1^3 + z_2z_6^2 - z_2z_3z_5 - 3z_1z_3z_4 + 2z_3^2z_6, \\
\mathbb{D}_9 = 2z_1^2z_3 + z_3^2z_4 - z_5^2z_6 + z_2z_4z_6, \\
\mathbb{D}_{10} = -z_1z_3^2 + z_2z_4^2 - 2z_4z_5^2 - z_1z_2z_6, \\
\mathbb{D}_{11} = -z_3^3 + z_1z_5^2 - z_1z_2z_4 - 3z_2z_3z_6 + 2z_2^2z_5, \\
\mathbb{D}_{12} = -z_1^2z_2 + z_3z_5^2 - 2z_5z_6^2 - z_2z_3z_4, \\
\mathbb{D}_\infty = z_4z_5z_6
\end{cases} \tag{1.11}$$

are the senary cubic forms (cubic forms in six variables).

Theorem 1.1. (Main Theorem 1). *The invariant decomposition formula for the simple group $PSL(2,13)$ of order 1092 is given as follows:*

$$\left[\frac{\eta^2(z)}{\eta^2(13z)}\right]^5 = \frac{\Phi_{12}(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))}{(a_1(z)a_2(z)a_3(z)a_4(z)a_5(z)a_6(z))^2}, \tag{1.12}$$

where $\Phi_{12}(z_1,z_2,z_3,z_4,z_5,z_6)$ is an invariant of degree 12 associated to $PSL(2,13)$, and $(z_1z_2z_3z_4z_5z_6)^2$ is invariant under the action of the image of a Borel subgroup of $PSL(2,13)$ (i.e. a maximal subgroup of order 78 of $PSL(2,13)$).

One of the significance of the invariant decomposition formulas (1.6), (1.7) and (1.12) comes from that the Hauptmodul $f(z) = (\eta(z)/\eta(pz))^{24/(p-1)}$ is a modular unit in the rational function field of $X_0(p)$ for $p = 5, 7, 13$. Its divisor is given by $\mathrm{div}(f) = \{(0) - (\infty)\}$. Hence, (1.6), (1.7) and (1.12) give some decomposition formulas of the modular units which play an important role both in the function field theory and applications to number theory (see [KL] for more details).

The other significance of the invariant decomposition formulas (1.6), (1.7) and (1.12) comes from the following fact: let H be a maximal subgroup of $PSL(2,p)$ for $p = 5$, 7 and 13 with order 10, 21 and 78, respectively. Then

$$[\Gamma : \Gamma_0(p)] = [PSL(2,p) : H] = p+1 \quad \text{for } p = 5,\ 7 \text{ and } 13, \text{ where } \Gamma = PSL(2,\mathbb{Z}).$$

Perhaps the most significance of the invariant decomposition formulas (1.6), (1.7) and (1.12) is the relation with the Monster simple group $\mathbb{M}$ and monstrous moonshine. In fact, elements of type $5B$, $7B$, $13B$ in the Monster correspond to elements in the conjugacy classes $5A$, $7A$, $13A$ of the Conway group Co_1 with cycle shapes 5^61^{-6}, 7^41^{-4}, 13^21^{-2}. The modular groups corresponding to these elements of the Monster are $\Gamma_0(5)$, $\Gamma_0(7)$ and $\Gamma_0(13)$ (see [Bor], [CN] and [CC]). Moreover, the Hauptmoduln $(\eta(z)/\eta(5z))^6$, $(\eta(z)/\eta(7z))^4$ and $(\eta(z)/\eta(13z))^2$ correspond exactly to the cycle shapes 5^61^{-6}, 7^41^{-4}, 13^21^{-2}, respectively (see Appendix for more details).

In [K1], Klein obtained the modular equation of degree 14 for $\Gamma_0(13)$ (see [CY], p.267 for the modular relation for $\Gamma_0(13)$). Combining with Theorem 1.1, we give a new expression of the classical j-function in terms of theta constants associated with $\Gamma(13)$.

Corollary 1.2. *The following formulas hold:*

$$j(z) = \frac{(\tau^2+5\tau+13)(\tau^4+247\tau^3+3380\tau^2+15379\tau+28561)^3}{\tau^{13}} \tag{1.13}$$

and

$$j(13z) = \frac{(\tau^2+5\tau+13)(\tau^4+7\tau^3+20\tau^2+19\tau+1)^3}{\tau}, \tag{1.14}$$

where the Hauptmodul

$$\tau = \left(\frac{\eta(z)}{\eta(13z)}\right)^2 = \sqrt[5]{\frac{\Phi_{12}(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z))}{(a_1(z)a_2(z)a_3(z)a_4(z)a_5(z)a_6(z))^2}}. \tag{1.15}$$

Note that in the right hand-side of modular equations (1.13) and (1.14), we have the following factorizations over $\mathbb{Q}(\sqrt{13})$:

$$\begin{aligned}&\tau^4 + 247\tau^3 + 3380\tau^2 + 15379\tau + 28561\\ &= \left(\tau^2 + \frac{247 + 65\sqrt{13}}{2}\tau + \frac{1859 + 507\sqrt{13}}{2}\right)\left(\tau^2 + \frac{247 - 65\sqrt{13}}{2}\tau + \frac{1859 - 507\sqrt{13}}{2}\right)\end{aligned}$$

and

$$\begin{aligned}&\tau^4 + 7\tau^3 + 20\tau^2 + 19\tau + 1\\ &= \left(\tau^2 + \frac{7 + \sqrt{13}}{2}\tau + \frac{11 + 3\sqrt{13}}{2}\right)\left(\tau^2 + \frac{7 - \sqrt{13}}{2}\tau + \frac{11 - 3\sqrt{13}}{2}\right).\end{aligned}$$

In [De1], Deligne gave a modular interpretation of why such factorizations exist. More generally, in [De2], Deligne showed that for $p = 3, 5, 7, 13$, the corresponding modular equations of degree 4, 6, 8, 14 have two relatively prime conjugate factors over some real quadratic fields. All of these factorizations have nice geometric interpretations. This is the other significance of the invariant decomposition formulas (1.6), (1.7) and (1.12) since they give the expressions of the Hauptmodul $\tau = (\eta(z)/\eta(pz))^{24/(p-1)}$ for $p = 5, 7, 13$.

In his notebooks (see [R5], p.326 and [R6], p.244), Ramanujan (see also [Ev] or [B1], p.373-375) obtained the following modular equation of degree 13:

$$1 + G_1(z)G_5(z) + G_2(z)G_3(z) + G_4(z)G_6(z) = -\frac{\eta^2(z/13)}{\eta^2(z)} \tag{1.16}$$

with

$$G_1(z)G_5(z)G_2(z)G_3(z)G_4(z)G_6(z) = -1, \tag{1.17}$$

where

$$G_m(z) := G_{m,p}(z) := (-1)^m q^{m(3m-p)/(2p^2)} \frac{f(-q^{2m/p}, -q^{1-2m/p})}{f(-q^{m/p}, -q^{1-m/p})}.$$

Here, m is a positive integer, $p = 13$, $q = e^{2\pi iz}$, and Ramanujan's general theta function $f(a, b)$ is given by

$$f(a, b) := \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2}, \quad |ab| < 1,$$

We call (1.16) the standard modular equation of degree 13. In contrast with it, we find the following invariant decomposition formula (exotic modular equation) which has the same form as (1.16), but with different kinds of modular parametrizations. Let us define the following senary quadratic forms (quadratic forms in six variables):

$$\begin{cases} \mathbb{A}_0 = z_1z_4 + z_2z_5 + z_3z_6, \\ \mathbb{A}_1 = z_1^2 - 2z_3z_4, \\ \mathbb{A}_2 = -z_5^2 - 2z_2z_4, \\ \mathbb{A}_3 = z_2^2 - 2z_1z_5, \\ \mathbb{A}_4 = z_3^2 - 2z_2z_6, \\ \mathbb{A}_5 = -z_4^2 - 2z_1z_6, \\ \mathbb{A}_6 = -z_6^2 - 2z_3z_5. \end{cases} \tag{1.18}$$

Theorem 1.3. (Main Theorem 2). *The following invariant decomposition formula (exotic modular equation) holds:*

$$\frac{\Psi_2(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z))}{\mathbb{A}_0(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z))^2} = 0, \tag{1.19}$$

where the quadric

$$\Psi_2(z_1, z_2, z_3, z_4, z_5, z_6) := \mathbb{A}_0^2 + \mathbb{A}_1\mathbb{A}_5 + \mathbb{A}_2\mathbb{A}_3 + \mathbb{A}_4\mathbb{A}_6 \tag{1.20}$$

is an invariant associated to $PSL(2, 13)$ and $\mathbb{A}_0^2$ is invariant under the action of the image of a Borel subgroup of $PSL(2, 13)$ (i.e. a maximal subgroup of order 78 of $PSL(2, 13)$).

Note that (1.19) has the same form as (1.16). However, in contrast with (1.17), it can be proved that (see section 3, (3.22))

$$\prod_{j=1}^{6} \frac{\mathbb{A}_j(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z))}{\mathbb{A}_0(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z))} \neq -1. \tag{1.21}$$

Moreover, the invariant quadric (1.20) is closely related to the exceptional Lie group G_2 (see [Y] for more details).

In [K], Klein investigated the modular equation of degree six in connection with the Jacobian equation of degree six, which can be used to solve the general quintic equation. In [K2] and [K3], he studied the modular equation of degree eight in connection with the Jacobian equation of degree eight, which can be used to solve the algebraic equation of degree eight with Galois group $PSL(2, 7)$. However, he did not investigate the last case, the modular equation of degree fourteen. In this paper, we find the

Jacobian equation of degree fourteen, which corresponds to the modular equation of degree fourteen. Surprisingly, we find the other exotic modular equation of degree fourteen, which does not correspond to the Jacobian equation. It plays an central role in the proof of Theorem 1.1. As an application, we obtain the following quartic four-fold $\Phi_4(z_1, z_2, z_3, z_4, z_5, z_6) = 0$, where

$$\Phi_4 := (z_3z_4^3+z_1z_5^3+z_2z_6^3)-(z_6z_1^3+z_4z_2^3+z_5z_3^3)+3(z_1z_2z_4z_5+z_2z_3z_5z_6+z_3z_1z_6z_4), \quad (1.22)$$

which is just the quadric (1.20) up to a constant. It is a higher-dimensional counterpart of the Klein quartic curve (see [K2]) and the Klein cubic threefold (see [K4]). Its significance comes from that it gives the geometry of the classical modular curve $X(13)$ of genus 50:

Corollary 1.4. *The coordinates* $(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z))$ *map* $X(13)$ *into the quartic four-fold* $\Phi_4(z_1, z_2, z_3, z_4, z_5, z_6) = 0$ *in* $\mathbb{CP}^5$.

This paper consists of four sections. In section two, we give a six-dimensional representation of $PSL(2, 13)$ defined over $\mathbb{Q}(e^{\frac{2\pi i}{13}})$ and the transformation formulas for theta constants associated with $\Gamma(13)$. In section three, we give a seven-dimensional representation of $PSL(2, 13)$ which is deduced from our six-dimensional representation. As an application, we obtain the exotic modular equation of degree 13 and give the geometry of the modular curve $X(13)$. Thus, we give the proof of Theorem 1.3. In section four, we give a fourteen-dimensional representation of $PSL(2, 13)$ which is also deduced from our six-dimensional representation. By this representation, we find the exotic modular equation of degree fourteen. Using it, we give the proof of Theorem 1.1.

Acknowedgements: I thank Pierre Deligne for his comments on an earlier version of this paper.

2. Six-dimensional representations of $PSL(2, 13)$ and transformation formulas for theta constants

Let us recall the Weil representation of $SL_2(\mathbb{F}_p)$ (see [W]) which is modelled on the space $L^2(\mathbb{F}_p)$ of all complex valued functions on $\mathbb{F}_p$ where p is a prime. Denote by $L^2(\mathbb{F}_p)$ the p-dimensional complex vector space of all (square-integrable) complex valued functions on $\mathbb{F}_p$ with respect to counting measure, that is, all functions from $\mathbb{F}_p$ to the complex numbers. We can decompose $L^2(\mathbb{F}_p)$ as the direct sum of the space V^+ of even functions and the space V^- of odd functions. The space V^- has dimension $(p-1)/2$ and its associated projective space $\mathbb{P}(V^-)$ has dimension $(p-3)/2$. In particular, for $p = 13$, the Weil representation for $SL(2, 13)$ is given as follows:

$$s = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad t = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad h = \begin{pmatrix} 7 & 0 \\ 0 & 2 \end{pmatrix}.$$

In this section, we will study the six-dimensional representation of the simple group $PSL(2,13)$ of order 1092, which acts on the five-dimensional projective space

$$\mathbb{P}^5 = \{(z_1, z_2, z_3, z_4, z_5, z_6) : z_i \in \mathbb{C} \quad (i = 1, 2, 3, 4, 5, 6)\}.$$

Let $\zeta = \exp(2\pi i/13)$ and

$$\begin{cases} \theta_1 = \zeta + \zeta^3 + \zeta^9, \\ \theta_2 = \zeta^2 + \zeta^6 + \zeta^5, \\ \theta_3 = \zeta^4 + \zeta^{12} + \zeta^{10}, \\ \theta_4 = \zeta^8 + \zeta^{11} + \zeta^7. \end{cases} \tag{2.1}$$

We find that

$$\begin{cases} \theta_1 + \theta_2 + \theta_3 + \theta_4 = -1, \\ \theta_1\theta_2 + \theta_1\theta_3 + \theta_1\theta_4 + \theta_2\theta_3 + \theta_2\theta_4 + \theta_3\theta_4 = 2, \\ \theta_1\theta_2\theta_3 + \theta_1\theta_2\theta_4 + \theta_1\theta_3\theta_4 + \theta_2\theta_3\theta_4 = 4, \\ \theta_1\theta_2\theta_3\theta_4 = 3. \end{cases}$$

Hence, θ_1, θ_2, θ_3 and θ_4 satisfy the quartic equation

$$z^4 + z^3 + 2z^2 - 4z + 3 = 0,$$

which can be decomposed as two quadratic equations

$$\left(z^2 + \frac{1+\sqrt{13}}{2}z + \frac{5+\sqrt{13}}{2}\right)\left(z^2 + \frac{1-\sqrt{13}}{2}z + \frac{5-\sqrt{13}}{2}\right) = 0$$

over the real quadratic field $\mathbb{Q}(\sqrt{13})$. Therefore, the four roots are given as follows:

$$z_{1,2} = \frac{1}{2}\left(-\frac{1+\sqrt{13}}{2} \pm \sqrt{\frac{-13-3\sqrt{13}}{2}}\right) = \frac{1}{4}\left(-1-\sqrt{13} \pm \sqrt{-26-6\sqrt{13}}\right),$$

$$z_{3,4} = \frac{1}{2}\left(-\frac{1-\sqrt{13}}{2} \pm \sqrt{\frac{-13+3\sqrt{13}}{2}}\right) = \frac{1}{4}\left(-1+\sqrt{13} \pm \sqrt{-26+6\sqrt{13}}\right).$$

Note that

$$\mathrm{Re}(\theta_1) = \cos\frac{2\pi}{13} + \cos\frac{6\pi}{13} - \cos\frac{5\pi}{13} > 0, \quad \mathrm{Im}(\theta_1) = \sin\frac{2\pi}{13} + \sin\frac{6\pi}{13} - \sin\frac{5\pi}{13} > 0.$$

We have

$$\theta_1 = \frac{1}{4}\left(-1+\sqrt{13}+\sqrt{-26+6\sqrt{13}}\right).$$

Similarly,

$$\mathrm{Re}(\theta_2)<0,\quad \mathrm{Im}(\theta_2)>0,\quad \theta_2=\frac{1}{4}\left(-1-\sqrt{13}+\sqrt{-26-6\sqrt{13}}\right).$$

$$\mathrm{Re}(\theta_3)>0,\quad \mathrm{Im}(\theta_3)<0,\quad \theta_3=\frac{1}{4}\left(-1+\sqrt{13}-\sqrt{-26+6\sqrt{13}}\right).$$

$$\mathrm{Re}(\theta_4)<0,\quad \mathrm{Im}(\theta_4)<0,\quad \theta_4=\frac{1}{4}\left(-1-\sqrt{13}-\sqrt{-26-6\sqrt{13}}\right).$$

Moreover, we find that

$$\begin{cases}\theta_1+\theta_3+\theta_2+\theta_4=-1,\\ \theta_1+\theta_3-\theta_2-\theta_4=\sqrt{13},\\ \theta_1-\theta_3-\theta_2+\theta_4=-\sqrt{-13+2\sqrt{13}},\\ \theta_1-\theta_3+\theta_2-\theta_4=\sqrt{-13-2\sqrt{13}}.\end{cases}\tag{2.2}$$

Let

$$M=\begin{pmatrix}\zeta-\zeta^{12} & \zeta^3-\zeta^{10} & \zeta^9-\zeta^4\\ \zeta^3-\zeta^{10} & \zeta^9-\zeta^4 & \zeta-\zeta^{12}\\ \zeta^9-\zeta^4 & \zeta-\zeta^{12} & \zeta^3-\zeta^{10}\end{pmatrix},\quad N=\begin{pmatrix}\zeta^5-\zeta^8 & \zeta^2-\zeta^{11} & \zeta^6-\zeta^7\\ \zeta^2-\zeta^{11} & \zeta^6-\zeta^7 & \zeta^5-\zeta^8\\ \zeta^6-\zeta^7 & \zeta^5-\zeta^8 & \zeta^2-\zeta^{11}\end{pmatrix}.\tag{2.3}$$

Then $MN=NM=-\sqrt{13}I$ and $M^2+N^2=-13I$. Put

$$S=-\frac{1}{\sqrt{13}}\begin{pmatrix}-M & N\\ N & M\end{pmatrix}.\tag{2.4}$$

Then $S^2=I$. In fact,

$$S=-\frac{2i}{\sqrt{13}}\begin{pmatrix}-\sin\frac{2\pi}{13} & -\sin\frac{6\pi}{13} & \sin\frac{5\pi}{13} & \sin\frac{3\pi}{13} & \sin\frac{4\pi}{13} & \sin\frac{\pi}{13}\\ -\sin\frac{6\pi}{13} & \sin\frac{5\pi}{13} & -\sin\frac{2\pi}{13} & \sin\frac{4\pi}{13} & \sin\frac{\pi}{13} & \sin\frac{3\pi}{13}\\ \sin\frac{5\pi}{13} & -\sin\frac{2\pi}{13} & -\sin\frac{6\pi}{13} & \sin\frac{\pi}{13} & \sin\frac{3\pi}{13} & \sin\frac{4\pi}{13}\\ \sin\frac{3\pi}{13} & \sin\frac{4\pi}{13} & \sin\frac{\pi}{13} & \sin\frac{2\pi}{13} & \sin\frac{6\pi}{13} & -\sin\frac{5\pi}{13}\\ \sin\frac{4\pi}{13} & \sin\frac{\pi}{13} & \sin\frac{3\pi}{13} & \sin\frac{6\pi}{13} & -\sin\frac{5\pi}{13} & \sin\frac{2\pi}{13}\\ \sin\frac{\pi}{13} & \sin\frac{3\pi}{13} & \sin\frac{4\pi}{13} & -\sin\frac{5\pi}{13} & \sin\frac{2\pi}{13} & \sin\frac{6\pi}{13}\end{pmatrix}.\tag{2.5}$$

Note that

$$\frac{\sin\frac{2\pi}{13}\sin\frac{5\pi}{13}\sin\frac{6\pi}{13}}{\sin\frac{\pi}{13}\sin\frac{3\pi}{13}\sin\frac{4\pi}{13}}=\frac{-\sqrt{\frac{-13-3\sqrt{13}}{2}}}{-\sqrt{\frac{-13+3\sqrt{13}}{2}}}=\frac{3+\sqrt{13}}{2},$$

which is a fundamental unit of $\mathbb{Q}(\sqrt{13})$. Let

$$T = \mathrm{diag}(\zeta^7, \zeta^{11}, \zeta^8, \zeta^6, \zeta^2, \zeta^5). \tag{2.6}$$

Theorem 2.1. *Let $G = \langle S, T\rangle$. Then $G \cong PSL(2, 13)$.*

Proof. We have

$$ST = -\frac{1}{\sqrt{13}} \begin{pmatrix} \zeta^6-\zeta^8 & \zeta^8-\zeta & \zeta^{12}-\zeta^4 & \zeta^{11}-\zeta & \zeta^4-1 & \zeta^{11}-\zeta^{12} \\ \zeta^4-\zeta^{10} & \zeta^2-\zeta^7 & \zeta^7-\zeta^9 & \zeta^8-\zeta^4 & \zeta^8-\zeta^9 & \zeta^{10}-1 \\ \zeta^{11}-\zeta^3 & \zeta^{10}-\zeta^{12} & \zeta^5-\zeta^{11} & \zeta^{12}-1 & \zeta^7-\zeta^{10} & \zeta^7-\zeta^3 \\ \zeta^{12}-\zeta^2 & 1-\zeta^9 & \zeta-\zeta^2 & \zeta^7-\zeta^5 & \zeta^5-\zeta^{12} & \zeta-\zeta^9 \\ \zeta^9-\zeta^5 & \zeta^4-\zeta^5 & 1-\zeta^3 & \zeta^9-\zeta^3 & \zeta^{11}-\zeta^6 & \zeta^6-\zeta^4 \\ 1-\zeta & \zeta^3-\zeta^6 & \zeta^{10}-\zeta^6 & \zeta^2-\zeta^{10} & \zeta^3-\zeta & \zeta^8-\zeta^2 \end{pmatrix}. \tag{2.7}$$

On the other hand,

$$\begin{aligned} (ST)^{-1} &= T^{-1}S \\ &= -\frac{1}{\sqrt{13}} \begin{pmatrix} \zeta^5-\zeta^7 & \zeta^3-\zeta^9 & \zeta^{10}-\zeta^2 & \zeta^{11}-\zeta & \zeta^8-\zeta^4 & \zeta^{12}-1 \\ \zeta^{12}-\zeta^5 & \zeta^6-\zeta^{11} & \zeta-\zeta^3 & \zeta^4-1 & \zeta^8-\zeta^9 & \zeta^7-\zeta^{10} \\ \zeta^9-\zeta & \zeta^4-\zeta^6 & \zeta^2-\zeta^8 & \zeta^{11}-\zeta^{12} & \zeta^{10}-1 & \zeta^7-\zeta^3 \\ \zeta^{12}-\zeta^2 & \zeta^9-\zeta^5 & 1-\zeta & \zeta^8-\zeta^6 & \zeta^{10}-\zeta^4 & \zeta^3-\zeta^{11} \\ 1-\zeta^9 & \zeta^4-\zeta^5 & \zeta^3-\zeta^6 & \zeta-\zeta^8 & \zeta^7-\zeta^2 & \zeta^{12}-\zeta^{10} \\ \zeta-\zeta^2 & 1-\zeta^3 & \zeta^{10}-\zeta^6 & \zeta^4-\zeta^{12} & \zeta^9-\zeta^7 & \zeta^{11}-\zeta^5 \end{pmatrix}. \end{aligned}$$

We will calculate $(ST)^2 = \frac{1}{13}(a_{ij})$. Without loss of generality, we will compute a_{1i}, $i = 1, 2, 3, 4, 5, 6$. Here,

$$a_{11} = -2\zeta - 2\zeta^2 + 2\zeta^3 - 2\zeta^9 + 2\zeta^{10} + 2\zeta^{11} - \zeta^5 + \zeta^7.$$

By the identity

$$\sqrt{13} = \zeta + \zeta^{12} + \zeta^3 + \zeta^{10} + \zeta^9 + \zeta^4 - \zeta^5 - \zeta^8 - \zeta^2 - \zeta^{11} - \zeta^6 - \zeta^7, \tag{2.8}$$

we have

$$(\zeta^5 - \zeta^7)\sqrt{13} = 2\zeta + 2\zeta^2 - 2\zeta^3 + 2\zeta^9 - 2\zeta^{10} - 2\zeta^{11} + \zeta^5 - \zeta^7.$$

Hence, $a_{11} = -(\zeta^5 - \zeta^7)\sqrt{13}$. Similarly,

$$\begin{aligned} a_{12} &= -2\zeta^2 - 2\zeta^4 + 2\zeta^5 - 2\zeta^7 + 2\zeta^8 + 2\zeta^{10} - \zeta^3 + \zeta^9 = -(\zeta^3 - \zeta^9)\sqrt{13}, \\ a_{13} &= -2 + 2\zeta^3 + 2\zeta^5 - 2\zeta^7 - 2\zeta^9 + 2\zeta^{12} + \zeta^2 - \zeta^{10} = -(\zeta^{10} - \zeta^2)\sqrt{13}, \\ a_{14} &= 2 + 2\zeta^4 + 2\zeta^5 - 2\zeta^7 - 2\zeta^8 - 2\zeta^{12} + \zeta^{11} - \zeta = -(\zeta^{11} - \zeta)\sqrt{13}, \\ a_{15} &= 2 + 2\zeta + 2\zeta^3 - 2\zeta^9 - 2\zeta^{11} - 2\zeta^{12} + \zeta^8 - \zeta^4 = -(\zeta^8 - \zeta^4)\sqrt{13}, \\ a_{16} &= 2\zeta - 2\zeta^2 + 2\zeta^4 - 2\zeta^8 + 2\zeta^{10} - 2\zeta^{11} + \zeta^{12} - 1 = -(\zeta^{12} - 1)\sqrt{13}. \end{aligned}$$

The other terms can be calculated in the same way. In conclusion, we find that $(ST)^2 = (ST)^{-1}$, i.e., $(ST)^3 = 1$. Hence, we have

$$S^2 = T^{13} = (ST)^3 = 1. \tag{2.9}$$

Let $u = ST$ and $v = S$. Then $uv = STS$. Hence,

$$u^3 = v^2 = (uv)^{13} = 1. \tag{2.10}$$

According to [S], put

$$(2, 3, n; p) := \langle u, v : u^3 = v^2 = (uv)^n = (u^{-1}v^{-1}uv)^p = 1\rangle.$$

Let $P = (uv)^{-1}$ and $Q = (uv)^2u$. Then $u = P^2Q$ and $v = P^3Q$. In [S], Sinkov proved the following:

Theorem 2.2. (see [S]). *Two operators of periods* 2 *and* 3 *generate the simple group of order* 1092 *if and only if they satisfy one of the following sets of independent relations:*

$$\begin{aligned} A: &\quad (2, 3, 7; 6),\\ B: &\quad (2, 3, 7; 7),\\ C: &\quad (2, 3, 7); (Q^2P^6)^3 = 1,\\ D: &\quad (2, 3, 13); (Q^3P^4)^3 = 1. \end{aligned}$$

In our case, $P = ST^{-1}S$ and $Q = ST^3$. We have

$$Q^3 = -\frac{1}{\sqrt{13}}\begin{pmatrix} \zeta^{11}-\zeta & \zeta^{12}-\zeta^8 & 1-\zeta & \zeta^6-\zeta^4 & \zeta^4-\zeta^{11} & 1-\zeta^8 \\ 1-\zeta^9 & \zeta^8-\zeta^9 & \zeta^4-\zeta^7 & 1-\zeta^7 & \zeta^2-\zeta^{10} & \zeta^{10}-\zeta^8 \\ \zeta^{10}-\zeta^{11} & 1-\zeta^3 & \zeta^7-\zeta^3 & \zeta^{12}-\zeta^7 & 1-\zeta^{11} & \zeta^5-\zeta^{12} \\ \zeta^9-\zeta^7 & \zeta^2-\zeta^9 & \zeta^5-1 & \zeta^2-\zeta^{12} & \zeta-\zeta^5 & 1-\zeta^{12} \\ \zeta^6-1 & \zeta^3-\zeta^{11} & \zeta^5-\zeta^3 & 1-\zeta^4 & \zeta^5-\zeta^4 & \zeta^9-\zeta^6 \\ \zeta^6-\zeta & \zeta^2-1 & \zeta-\zeta^8 & \zeta^3-\zeta^2 & 1-\zeta^{10} & \zeta^6-\zeta^{10} \end{pmatrix}.$$

$$P^4 = -\frac{1}{\sqrt{13}}\begin{pmatrix} \zeta^7-1 & \zeta^2-\zeta^7 & \zeta^6-\zeta^8 & \zeta^2-\zeta^{11} & \zeta^5-\zeta^6 & \zeta^8-\zeta^{11} \\ \zeta^2-\zeta^7 & \zeta^{11}-1 & \zeta^5-\zeta^{11} & \zeta^7-\zeta^8 & \zeta^5-\zeta^8 & \zeta^6-\zeta^2 \\ \zeta^6-\zeta^8 & \zeta^5-\zeta^{11} & \zeta^8-1 & \zeta^2-\zeta^5 & \zeta^{11}-\zeta^7 & \zeta^6-\zeta^7 \\ \zeta^2-\zeta^{11} & \zeta^7-\zeta^8 & \zeta^2-\zeta^5 & \zeta^6-1 & \zeta^{11}-\zeta^6 & \zeta^7-\zeta^5 \\ \zeta^5-\zeta^6 & \zeta^5-\zeta^8 & \zeta^{11}-\zeta^7 & \zeta^{11}-\zeta^6 & \zeta^2-1 & \zeta^8-\zeta^2 \\ \zeta^8-\zeta^{11} & \zeta^6-\zeta^2 & \zeta^6-\zeta^7 & \zeta^7-\zeta^5 & \zeta^8-\zeta^2 & \zeta^5-1 \end{pmatrix}.$$

$$Q^3P^4 = -\frac{1}{\sqrt{13}}\begin{pmatrix} \zeta^7-\zeta^5 & \zeta^2-\zeta^9 & \zeta^{10}-\zeta^5 & \zeta^6-\zeta^3 & \zeta^3-\zeta^7 & \zeta^{10}-\zeta^9 \\ \zeta^{12}-\zeta^6 & \zeta^{11}-\zeta^6 & \zeta^5-\zeta^3 & \zeta^{12}-\zeta^3 & \zeta^2-\zeta & \zeta-\zeta^{11} \\ \zeta^6-\zeta & \zeta^4-\zeta^2 & \zeta^8-\zeta^2 & \zeta^9-\zeta^8 & \zeta^4-\zeta & \zeta^5-\zeta^9 \\ \zeta^{10}-\zeta^7 & \zeta^6-\zeta^{10} & \zeta^4-\zeta^3 & \zeta^6-\zeta^8 & \zeta^{11}-\zeta^4 & \zeta^3-\zeta^8 \\ \zeta^{10}-\zeta & \zeta^{12}-\zeta^{11} & \zeta^2-\zeta^{12} & \zeta-\zeta^7 & \zeta^2-\zeta^7 & \zeta^8-\zeta^{10} \\ \zeta^5-\zeta^4 & \zeta^{12}-\zeta^9 & \zeta^4-\zeta^8 & \zeta^7-\zeta^{12} & \zeta^9-\zeta^{11} & \zeta^5-\zeta^{11} \end{pmatrix},$$

$$(Q^3P^4)^2 = -\frac{1}{\sqrt{13}}\begin{pmatrix} \zeta^8-\zeta^6 & \zeta^7-\zeta & \zeta^{12}-\zeta^7 & \zeta^6-\zeta^3 & \zeta^{12}-\zeta^3 & \zeta^9-\zeta^8 \\ \zeta^4-\zeta^{11} & \zeta^7-\zeta^2 & \zeta^{11}-\zeta^9 & \zeta^3-\zeta^7 & \zeta^2-\zeta & \zeta^4-\zeta \\ \zeta^8-\zeta^3 & \zeta^{10}-\zeta^8 & \zeta^{11}-\zeta^5 & \zeta^{10}-\zeta^9 & \zeta-\zeta^{11} & \zeta^5-\zeta^9 \\ \zeta^{10}-\zeta^7 & \zeta^{10}-\zeta & \zeta^5-\zeta^4 & \zeta^5-\zeta^7 & \zeta^6-\zeta^{12} & \zeta-\zeta^6 \\ \zeta^6-\zeta^{10} & \zeta^{12}-\zeta^{11} & \zeta^{12}-\zeta^9 & \zeta^9-\zeta^2 & \zeta^6-\zeta^{11} & \zeta^2-\zeta^4 \\ \zeta^4-\zeta^3 & \zeta^2-\zeta^{12} & \zeta^4-\zeta^8 & \zeta^5-\zeta^{10} & \zeta^3-\zeta^5 & \zeta^2-\zeta^8 \end{pmatrix},$$

and $(Q^3P^4)^3 = -I$. Note that in the projective coordinates, this means that $(Q^3P^4)^3 = 1$. Hence, we prove that the elements u and v above satisfy the following relations: $(2, 3, 13)$ and $(Q^3P^4)^3 = 1$, which is a presentation for the simple group $PSL(2, 13)$ of order 1092 by Theorem 3.2. Since the group is simple and the generating matrices are non-trivial, we must have $G = \langle u, v\rangle \cong PSL(2, 13)$. Hence, $\langle P, Q\rangle = \langle S, T\rangle = G$. This completes the proof of Theorem 2.1.

□

Recall that the theta functions with characteristic $\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} \in \mathbb{R}^2$ is defined by the following series which converges uniformly and absolutely on compact subsets of $\mathbb{C} \times \mathbb{H}$ (see [FK], p. 73):

$$\theta\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}(z, \tau) = \sum_{n\in\mathbb{Z}} \exp\left\{2\pi i\left[\frac{1}{2}\left(n+\frac{\epsilon}{2}\right)^2\tau + \left(n+\frac{\epsilon}{2}\right)\left(z+\frac{\epsilon'}{2}\right)\right]\right\}.$$

The modified theta constants is defined by (see [FK], p. 215)

$$\varphi_l(\tau) = \theta[\chi_l](0, k\tau),$$

where the characteristic $\chi_l = \begin{bmatrix} \frac{2l+1}{k} \\ 1 \end{bmatrix}$, $l = 0, \cdots, \frac{k-3}{2}$, for odd k and $\chi_l = \begin{bmatrix} \frac{2l}{k} \\ 0 \end{bmatrix}$, $l = 0, \cdots, \frac{k}{2}$, for even k. We have the following:

Theorem 2.3. (see [FK], p. 236). *For each odd integer $k \geq 5$, the map*

$$\Phi : \tau \mapsto (\varphi_0(\tau), \varphi_1(\tau), \cdots, \varphi_{\frac{k-5}{2}}(\tau), \varphi_{\frac{k-3}{2}}(\tau))$$

from $\mathbb{H} \cup \mathbb{Q} \cup \{\infty\}$ to $\mathbb{C}^{\frac{k-1}{2}}$, defines a holomorphic mapping from $\overline{\mathbb{H}/\Gamma(k)}$ into $\mathbb{CP}^{\frac{k-3}{2}}$.

In our case, $k = 13$, the map

$$\Phi : \tau \mapsto (\varphi_0(\tau), \varphi_1(\tau), \varphi_2(\tau), \varphi_3(\tau), \varphi_4(\tau), \varphi_5(\tau))$$

gives a holomorphic mapping from the modular curve $X(13) = \overline{\mathbb{H}/\Gamma(13)}$ into $\mathbb{CP}^5$, which corresponds to our six-dimensional representation, i.e., up to the constants, $z_1, \cdots, z_6$ are just modular forms $\varphi_0(\tau), \cdots, \varphi_5(\tau)$.

Let

$$\begin{cases} a_1(z) := e^{-\frac{11\pi i}{26}}\theta\left[\begin{smallmatrix}\frac{11}{13}\\ 1\end{smallmatrix}\right](0,13z) = q^{\frac{121}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+11n)}, \\ a_2(z) := e^{-\frac{7\pi i}{26}}\theta\left[\begin{smallmatrix}\frac{7}{13}\\ 1\end{smallmatrix}\right](0,13z) = q^{\frac{49}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+7n)}, \\ a_3(z) := e^{-\frac{5\pi i}{26}}\theta\left[\begin{smallmatrix}\frac{5}{13}\\ 1\end{smallmatrix}\right](0,13z) = q^{\frac{25}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+5n)}, \\ a_4(z) := -e^{-\frac{3\pi i}{26}}\theta\left[\begin{smallmatrix}\frac{3}{13}\\ 1\end{smallmatrix}\right](0,13z) = -q^{\frac{9}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+3n)}, \\ a_5(z) := e^{-\frac{9\pi i}{26}}\theta\left[\begin{smallmatrix}\frac{9}{13}\\ 1\end{smallmatrix}\right](0,13z) = q^{\frac{81}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+9n)}, \\ a_6(z) := e^{-\frac{\pi i}{26}}\theta\left[\begin{smallmatrix}\frac{1}{13}\\ 1\end{smallmatrix}\right](0,13z) = q^{\frac{1}{104}}\sum_{n\in\mathbb{Z}}(-1)^n q^{\frac{1}{2}(13n^2+n)}. \end{cases} \tag{2.11}$$

We use the standard notation $(a)_\infty := (a;q) := \prod_{k=0}^{\infty}(1-aq^k)$ for $a \in \mathbb{C}^\times$ and $|q|<1$ so that $\eta(z) = q^{1/24}(q;q)$. By the Jacobi triple product identity, we have

$$\theta\left[\begin{smallmatrix}\frac{l}{k}\\ 1\end{smallmatrix}\right](0,kz) = \exp\left(\frac{\pi i l}{2k}\right) q^{\frac{l^2}{8k}}(q^{\frac{k-l}{2}};q^k)(q^{\frac{k+l}{2}};q^k)(q^k;q^k),$$

where k is odd and $l = 1,3,5,\cdots,k-2$. Hence,

$$\begin{cases} a_1(z) = q^{\frac{121}{104}}(q;q^{13})(q^{12};q^{13})(q^{13};q^{13}), \\ a_2(z) = q^{\frac{49}{104}}(q^3;q^{13})(q^{10};q^{13})(q^{13};q^{13}), \\ a_3(z) = q^{\frac{25}{104}}(q^9;q^{13})(q^4;q^{13})(q^{13};q^{13}), \\ a_4(z) = -q^{\frac{9}{104}}(q^5;q^{13})(q^8;q^{13})(q^{13};q^{13}), \\ a_5(z) = q^{\frac{81}{104}}(q^2;q^{13})(q^{11};q^{13})(q^{13};q^{13}), \\ a_6(z) = q^{\frac{1}{104}}(q^6;q^{13})(q^7;q^{13})(q^{13};q^{13}). \end{cases} \tag{2.12}$$

Ramanujan's general theta function $f(a,b)$ is given by

$$f(a,b) := \sum_{n=-\infty}^{\infty} a^{n(n+1)/2}b^{n(n-1)/2}, \quad |ab|<1,$$

and

$$f(-q) := f(-q,-q^2) = \sum_{n=-\infty}^{\infty}(-1)^n q^{n(3n-1)/2} = (q;q)_\infty.$$

In his notebooks (see [R5], p.326, [R6], p.244 and [B1], p.372), Ramanujan obtained his modular equations of degree 13.

Theorem 2.4. *Define*

$$\mu_1 = \frac{f(-q^4,-q^9)}{q^{7/13}f(-q^2,-q^{11})}, \quad \mu_2 = \frac{f(-q^6,-q^7)}{q^{6/13}f(-q^3,-q^{10})},$$

$$\mu_3 = \frac{f(-q^2,-q^{11})}{q^{5/13}f(-q,-q^{12})}, \quad \mu_4 = \frac{f(-q^5,-q^8)}{q^{2/13}f(-q^4,-q^9)},$$

$$\mu_5 = \frac{q^{5/13}f(-q^3,-q^{10})}{f(-q^5,-q^8)}, \quad \mu_6 = \frac{q^{15/13}f(-q,-q^{12})}{f(-q^6,-q^7)}.$$

Then

$$1 + \frac{f^2(-q)}{qf^2(-q^{13})} = \mu_1\mu_2 - \mu_3\mu_5 - \mu_4\mu_6, \tag{2.13}$$

$$-4 - \frac{f^2(-q)}{qf^2(-q^{13})} = \frac{1}{\mu_1\mu_2} - \frac{1}{\mu_3\mu_5} - \frac{1}{\mu_4\mu_6}, \tag{2.14}$$

$$3 + \frac{f^2(-q)}{qf^2(-q^{13})} = \mu_2\mu_3\mu_4 - \mu_1\mu_5\mu_6, \tag{2.15}$$

where $\mu_1\mu_2\mu_3\mu_4\mu_5\mu_6 = 1$.

Let

$$G_m(z) := G_{m,p}(z) := (-1)^m q^{m(3m-p)/(2p^2)} \frac{f(-q^{2m/p},-q^{1-2m/p})}{f(-q^{m/p},-q^{1-m/p})}$$

where m is a positive integer, $p = 13$ and $q = e^{2\pi iz}$. Then the above three formulas (2.13), (2.14) and (2.15) are equivalent to the following (see [Ev] or [B1], p.373-375)

$$1 + G_1(z)G_5(z) + G_2(z)G_3(z) + G_4(z)G_6(z) = -\frac{\eta^2(z/13)}{\eta^2(z)}, \tag{2.16}$$

$$\frac{1}{G_1(z)G_5(z)} + \frac{1}{G_2(z)G_3(z)} + \frac{1}{G_4(z)G_6(z)} = 4 + \frac{\eta^2(z/13)}{\eta^2(z)}, \tag{2.17}$$

$$G_1(z)G_3(z)G_4(z) - \frac{1}{G_1(z)G_3(z)G_4(z)} = 3 + \frac{\eta^2(z/13)}{\eta^2(z)}, \tag{2.18}$$

where $G_1(z)G_2(z)G_3(z)G_4(z)G_5(z)G_6(z) = -1$. Moreover, there is the following formula (see [Ev] or [B1], p.375-376): for $t = q^{1/13}$,

$$\frac{1}{(t^2)_\infty(t^3)_\infty(t^{10})_\infty(t^{11})_\infty} + \frac{t}{(t^4)_\infty(t^6)_\infty(t^7)_\infty(t^9)_\infty} = \frac{1}{(t)_\infty(t^5)_\infty(t^8)_\infty(t^{12})_\infty}, \tag{2.19}$$

which is equivalent to, for $p = 13$, $G_1^{-1}(z)G_5^{-1}(z) + G_4(z)G_6(z) = 1$.

It is known that (see [Ev])

$$G(m; z) = (-1)^m F(2m/p; z)/F(m/p; z),$$

where

$$\begin{aligned} F(u; z) &= -i \sum_{k=-\infty}^{\infty} (-1)^k q^{(k+u+1/2)^2/2} \\ &= -iq^{(u+1/2)^2/2} \prod_{m=1}^{\infty} (1 - q^{m+u})(1 - q^{m-1-u})(1 - q^m) \end{aligned}$$

satisfies that $F(u + 1; z) = -F(u; z)$ and $F(-u; z) = -F(u; z)$. We have

$$G_1(z)G_5(z) = \frac{F(\frac{2}{13}; z)F(\frac{3}{13}; z)}{F(\frac{1}{13}; z)F(\frac{5}{13}; z)},$$

where

$$F\left(\frac{1}{13}; z\right) = -iq^{(\frac{1}{13}+\frac{1}{2})^2/2} \prod_{m=1}^{\infty} (1 - q^{m+\frac{1}{13}})(1 - q^{m-1-\frac{1}{13}})(1 - q^m),$$

$$F\left(\frac{5}{13}; z\right) = -iq^{(\frac{5}{13}+\frac{1}{2})^2/2} \prod_{m=1}^{\infty} (1 - q^{m+\frac{5}{13}})(1 - q^{m-1-\frac{5}{13}})(1 - q^m),$$

$$F\left(\frac{2}{13}; z\right) = -iq^{(\frac{2}{13}+\frac{1}{2})^2/2} \prod_{m=1}^{\infty} (1 - q^{m+\frac{2}{13}})(1 - q^{m-1-\frac{2}{13}})(1 - q^m),$$

$$F\left(\frac{3}{13}; z\right) = -iq^{(\frac{3}{13}+\frac{1}{2})^2/2} \prod_{m=1}^{\infty} (1 - q^{m+\frac{3}{13}})(1 - q^{m-1-\frac{3}{13}})(1 - q^m).$$

Note that

$$\prod_{m=1}^{\infty} (1 - q^{m+\frac{1}{13}}) = \frac{1}{1-t}(t)_\infty, \quad \prod_{m=1}^{\infty} (1 - q^{m-1-\frac{1}{13}}) = \frac{t-1}{t}(t^{12})_\infty,$$

$$\prod_{m=1}^{\infty} (1 - q^{m+\frac{5}{13}}) = \frac{1}{1-t^5}(t^5)_\infty, \quad \prod_{m=1}^{\infty} (1 - q^{m-1-\frac{5}{13}}) = \frac{t^5-1}{t^5}(t^8)_\infty,$$

$$\prod_{m=1}^{\infty} (1 - q^{m+\frac{2}{13}}) = \frac{1}{1-t^2}(t^2)_\infty, \quad \prod_{m=1}^{\infty} (1 - q^{m-1-\frac{2}{13}}) = \frac{t^2-1}{t^2}(t^{11})_\infty,$$

$$\prod_{m=1}^{\infty}(1-q^{m+\frac{3}{13}})=\frac{1}{1-t^3}(t^3)_\infty,\quad \prod_{m=1}^{\infty}(1-q^{m-1-\frac{3}{13}})=\frac{t^3-1}{t^3}(t^{10})_\infty,$$

where $(x)_\infty=\prod_{m=0}^{\infty}(1-xq^m)$ and $t=q^{\frac{1}{13}}$. Thus,

$$G_1(z)G_5(z)=\frac{(t^3)_\infty(t^{10})_\infty(t^2)_\infty(t^{11})_\infty}{(t)_\infty(t^{12})_\infty(t^5)_\infty(t^8)_\infty}.$$

Similarly,

$$G_2(z)G_3(z)=-\frac{(t^9)_\infty(t^4)_\infty(t^6)_\infty(t^7)_\infty}{t(t^3)_\infty(t^{10})_\infty(t^2)_\infty(t^{11})_\infty},\quad G_4(z)G_6(z)=\frac{t(t)_\infty(t^{12})_\infty(t^5)_\infty(t^8)_\infty}{(t^9)_\infty(t^4)_\infty(t^6)_\infty(t^7)_\infty}.$$

Note that $(t^k)_\infty=(t^k;t^{13})$ for $1\le k\le 13$. The above formula (2.16) is equivalent to

$$\begin{aligned}
&t(t^3;t^{13})^2(t^{10};t^{13})^2(t^2;t^{13})^2(t^{11};t^{13})^2(t^9;t^{13})(t^4;t^{13})(t^6;t^{13})(t^7;t^{13})\\
&-(t^9;t^{13})^2(t^4;t^{13})^2(t^6;t^{13})^2(t^7;t^{13})^2(t;t^{13})(t^{12};t^{13})(t^5;t^{13})(t^8;t^{13})\\
&+t^2(t;t^{13})^2(t^{12};t^{13})^2(t^5;t^{13})^2(t^8;t^{13})^2(t^3;t^{13})(t^{10};t^{13})(t^2;t^{13})(t^{11};t^{13})\\
&=[-1-\eta^2(z/13)/\eta^2(z)]t(t;t^{13})(t^2;t^{13})\cdots(t^{12};t^{13}),\quad t=q^{1/13}=e^{\frac{2\pi iz}{13}}.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
a_2(z)^2a_5(z)^2a_3(z)a_6(z)=&q^{\frac{11}{4}}(q^3;q^{13})^2(q^{10};q^{13})^2(q^2;q^{13})^2(q^{11};q^{13})^2\\
&\times(q^9;q^{13})(q^4;q^{13})(q^6;q^{13})(q^7;q^{13})(q^{13};q^{13})^6,\\
a_3(z)^2a_6(z)^2a_1(z)a_4(z)=&-q^{\frac{7}{4}}(q^9;q^{13})^2(q^4;q^{13})^2(q^6;q^{13})^2(q^7;q^{13})^2\\
&\times(q;q^{13})(q^{12};q^{13})(q^5;q^{13})(q^8;q^{13})(q^{13};q^{13})^6,\\
a_1(z)^2a_4(z)^2a_2(z)a_5(z)=&q^{\frac{15}{4}}(q;q^{13})^2(q^{12};q^{13})^2(q^5;q^{13})^2(q^8;q^{13})^2\\
&\times(q^3;q^{13})(q^{10};q^{13})(q^2;q^{13})(q^{11};q^{13})(q^{13};q^{13})^6,
\end{aligned}$$

This implies that,

$$\begin{aligned}
&a_1(z)^2a_4(z)^2a_2(z)a_5(z)+a_2(z)^2a_5(z)^2a_3(z)a_6(z)+a_3(z)^2a_6(z)^2a_1(z)a_4(z)\\
&=[-1-\eta^2(z)/\eta^2(13z)]\eta(z)\eta(13z)^5.
\end{aligned}\tag{2.20}$$

Similarly, the other three formulas (2.17), (2.18) and (2.19) are equivalent to the following:

$$\begin{aligned}
&a_1(z)^2a_4(z)^2a_3(z)a_6(z)+a_2(z)^2a_5(z)^2a_1(z)a_4(z)+a_3(z)^2a_6(z)^2a_2(z)a_5(z)\\
&=[4+\eta^2(z)/\eta^2(13z)]\eta(z)\eta(13z)^5.
\end{aligned}\tag{2.21}$$

$$a_4(z)^2a_5(z)^2a_6(z)^2 - a_1(z)^2a_2(z)^2a_3(z)^2 = [3 + \eta^2(z)/\eta^2(13z)]\eta(z)\eta(13z)^5, \quad (2.22)$$

and

$$\frac{1}{a_1(z)a_4(z)} + \frac{1}{a_2(z)a_5(z)} + \frac{1}{a_3(z)a_6(z)} = 0, \quad (2.23)$$

where $a_1(z)a_2(z)a_3(z)a_4(z)a_5(z)a_6(z) = -\eta(z)\eta(13z)^5$.

Let $\mathbf{A} = (a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z))^T$. The significance of our six dimensional representation of $PSL(2, 13)$ comes from the following:

Proposition 2.5. *If $z \in \mathbb{H}$, then the following relations hold:*

$$\mathbf{A}(z+1) = e^{-\frac{3\pi i}{4}} T\mathbf{A}(z), \quad \mathbf{A}\left(-\frac{1}{z}\right) = e^{\frac{\pi i}{4}}\sqrt{z}S\mathbf{A}(z), \quad (2.24)$$

where

$$T = \begin{pmatrix} \zeta^7 & & & & & \\ & \zeta^{11} & & & & \\ & & \zeta^8 & & & \\ & & & \zeta^6 & & \\ & & & & \zeta^2 & \\ & & & & & \zeta^5 \end{pmatrix},$$

$$S = -\frac{1}{\sqrt{13}} \begin{pmatrix} \zeta^{12}-\zeta & \zeta^{10}-\zeta^3 & \zeta^4-\zeta^9 & \zeta^5-\zeta^8 & \zeta^2-\zeta^{11} & \zeta^6-\zeta^7 \\ \zeta^{10}-\zeta^3 & \zeta^4-\zeta^9 & \zeta^{12}-\zeta & \zeta^2-\zeta^{11} & \zeta^6-\zeta^7 & \zeta^5-\zeta^8 \\ \zeta^4-\zeta^9 & \zeta^{12}-\zeta & \zeta^{10}-\zeta^3 & \zeta^6-\zeta^7 & \zeta^5-\zeta^8 & \zeta^2-\zeta^{11} \\ \zeta^5-\zeta^8 & \zeta^2-\zeta^{11} & \zeta^6-\zeta^7 & \zeta-\zeta^{12} & \zeta^3-\zeta^{10} & \zeta^9-\zeta^4 \\ \zeta^2-\zeta^{11} & \zeta^6-\zeta^7 & \zeta^5-\zeta^8 & \zeta^3-\zeta^{10} & \zeta^9-\zeta^4 & \zeta-\zeta^{12} \\ \zeta^6-\zeta^7 & \zeta^5-\zeta^8 & \zeta^2-\zeta^{11} & \zeta^9-\zeta^4 & \zeta-\zeta^{12} & \zeta^3-\zeta^{10} \end{pmatrix},$$

and $0 < arg\sqrt{z} \leq \pi/2$.

Proof. Recall that the following transformation formulas for theta constants (see [FK], p.216-217):

$$\theta\begin{bmatrix} \frac{2l+1}{k} \\ 1 \end{bmatrix}(0, k(z+1)) = \exp\left(\frac{\pi i}{4k}\right)\exp\left(\frac{\pi i}{k}(l^2+l)\right)\theta\begin{bmatrix} \frac{2l+1}{k} \\ 1 \end{bmatrix}(0, kz)$$

and

$$\theta\begin{bmatrix} \frac{2l+1}{k} \\ 1 \end{bmatrix}(0, k(-1/z)) = \frac{\sqrt{z}}{\sqrt{ik}}\exp\left(\frac{\pi i}{2k}\right) \times$$
$$\sum_{j=0}^{\frac{k-3}{2}} \left[\exp\left(2\pi i\frac{l(j+1)}{k}\right) + \exp\left(-\frac{\pi i}{k}\right)\exp\left(-2\pi i\frac{(l+1)j}{k}\right)\right]\theta\begin{bmatrix} \frac{2j+1}{k} \\ 1 \end{bmatrix}(0, kz),$$

where k is odd and $l = 0, 1, \cdots, \frac{k-3}{2}$. In our case, $k = 13$. By the first formula, we have

$$l = 5, e^{\frac{\pi i}{52}} e^{\frac{30\pi i}{13}} = e^{-\frac{3\pi i}{4}} \zeta^7; \quad l = 3, e^{\frac{\pi i}{52}} e^{\frac{12\pi i}{13}} = e^{-\frac{3\pi i}{4}} \zeta^{11}; \quad l = 2, e^{\frac{\pi i}{52}} e^{\frac{6\pi i}{13}} = e^{-\frac{3\pi i}{4}} \zeta^8.$$

$$l = 1, e^{\frac{\pi i}{52}} e^{\frac{2\pi i}{13}} = e^{-\frac{3\pi i}{4}} \zeta^6; \quad l = 4, e^{\frac{\pi i}{52}} e^{\frac{20\pi i}{13}} = e^{-\frac{3\pi i}{4}} \zeta^2; \quad l = 0, e^{\frac{\pi i}{52}} = e^{-\frac{3\pi i}{4}} \zeta^5.$$

This gives that $\mathbf{A}(z+1) = e^{-\frac{3\pi i}{4}} T\mathbf{A}(z)$.

By the second formula, for $l = 5$, we have

$$\theta \begin{bmatrix} \frac{11}{13} \\ 1 \end{bmatrix} (0, -13/z) = \frac{\sqrt{z}}{\sqrt{13i}} e^{\frac{\pi i}{26}} \sum_{j=0}^{5} \left[e^{\frac{2\pi i \cdot 5(j+1)}{13}} + e^{-\frac{\pi i}{13}} e^{-\frac{2\pi i \cdot 6j}{13}} \right] \theta \begin{bmatrix} \frac{2j+1}{13} \\ 1 \end{bmatrix} (0, 13z).$$

where

$$\frac{1}{\sqrt{13i}} e^{\frac{\pi i}{26}} = \frac{1}{\sqrt{13}} e^{-\frac{\pi i}{4}} e^{\frac{\pi i}{26}} = \frac{-1}{\sqrt{13}} e^{\frac{3\pi i}{4}} e^{\frac{\pi i}{26}} = \frac{-1}{\sqrt{13}} e^{\frac{\pi i}{4}} e^{\frac{7\pi i}{13}}.$$

Hence,

$$\begin{aligned}
&\theta \begin{bmatrix} \frac{11}{13} \\ 1 \end{bmatrix} (0, -13/z) \\
=&\frac{-1}{\sqrt{13}} \sqrt{z} e^{\frac{\pi i}{4}} e^{\frac{7\pi i}{13}} [(e^{\frac{10\pi i}{13}} + e^{-\frac{\pi i}{13}}) \theta \begin{bmatrix} \frac{1}{13} \\ 1 \end{bmatrix} (0, 13z) + (e^{\frac{20\pi i}{13}} + e^{-\pi i}) \theta \begin{bmatrix} \frac{3}{13} \\ 1 \end{bmatrix} (0, 13z) + \\
&+ (e^{\frac{30\pi i}{13}} + e^{-\frac{25\pi i}{13}}) \theta \begin{bmatrix} \frac{5}{13} \\ 1 \end{bmatrix} (0, 13z) + (e^{\frac{40\pi i}{13}} + e^{-\frac{37\pi i}{13}}) \theta \begin{bmatrix} \frac{7}{13} \\ 1 \end{bmatrix} (0, 13z) + \\
&+ (e^{\frac{50\pi i}{13}} + e^{-\frac{49\pi i}{13}}) \theta \begin{bmatrix} \frac{9}{13} \\ 1 \end{bmatrix} (0, 13z) + (e^{\frac{60\pi i}{13}} + e^{-\frac{61\pi i}{13}}) \theta \begin{bmatrix} \frac{11}{13} \\ 1 \end{bmatrix} (0, 13z)].
\end{aligned}$$

Here,

$$e^{\frac{7\pi i}{13}} (e^{\frac{10\pi i}{13}} + e^{-\frac{\pi i}{13}}) = e^{\frac{5\pi i}{13}} (\zeta^6 - \zeta^7), \quad e^{\frac{7\pi i}{13}} (e^{\frac{20\pi i}{13}} + e^{-\pi i}) = -e^{\frac{4\pi i}{13}} (\zeta^5 - \zeta^8),$$

$$e^{\frac{7\pi i}{13}} (e^{\frac{30\pi i}{13}} + e^{-\frac{25\pi i}{13}}) = e^{\frac{3\pi i}{13}} (\zeta^4 - \zeta^9), \quad e^{\frac{7\pi i}{13}} (e^{\frac{40\pi i}{13}} + e^{-\frac{37\pi i}{13}}) = e^{\frac{2\pi i}{13}} (\zeta^{10} - \zeta^3),$$

$$e^{\frac{7\pi i}{13}} (e^{\frac{50\pi i}{13}} + e^{-\frac{49\pi i}{13}}) = e^{\frac{\pi i}{13}} (\zeta^2 - \zeta^{11}), \quad e^{\frac{7\pi i}{13}} (e^{\frac{60\pi i}{13}} + e^{-\frac{61\pi i}{13}}) = \zeta^{12} - \zeta.$$

This implies that

$$\begin{aligned}
a_1 \left(-\frac{1}{z} \right) =& \frac{-1}{\sqrt{13}} \sqrt{z} e^{\frac{\pi i}{4}} [(\zeta^{12} - \zeta) a_1(z) + (\zeta^{10} - \zeta^3) a_2(z) + (\zeta^4 - \zeta^9) a_3(z) + \\
&+ (\zeta^5 - \zeta^8) a_4(z) + (\zeta^2 - \zeta^{11}) a_5(z) + (\zeta^6 - \zeta^7) a_6(z)].
\end{aligned}$$

For $l = 3, 2, 1, 4, 0$, we can prove the similar formulas. This gives that $\mathbf{A}\left(-\frac{1}{z}\right) = e^{\frac{\pi i}{4}} \sqrt{z} S\mathbf{A}(z)$. This completes the proof of Proposition 2.5.

□

3. Seven-dimensional representations of $PSL(2,13)$, exotic modular equation and geometry of modular curve $X(13)$

We will construct a seven-dimensional representation of $PSL(2,13)$ which is deduced from our six-dimensional representation. Let us study the action of ST^ν on the five dimensional projective space $\mathbb{P}^5=\{(z_1,z_2,z_3,z_4,z_5,z_6)\}$, where $\nu=0,1,\cdots,12$. Put

$$\alpha=\zeta+\zeta^{12}-\zeta^5-\zeta^8,\quad \beta=\zeta^3+\zeta^{10}-\zeta^2-\zeta^{11},\quad \gamma=\zeta^9+\zeta^4-\zeta^6-\zeta^7.$$

We find that

$$\begin{aligned}
&13ST^\nu(z_1)\cdot ST^\nu(z_4)\\
=&\beta z_1z_4+\gamma z_2z_5+\alpha z_3z_6+\\
&+\gamma\zeta^\nu z_1^2+\alpha\zeta^{9\nu}z_2^2+\beta\zeta^{3\nu}z_3^2-\gamma\zeta^{12\nu}z_4^2-\alpha\zeta^{4\nu}z_5^2-\beta\zeta^{10\nu}z_6^2+\\
&+(\alpha-\beta)\zeta^{5\nu}z_1z_2+(\beta-\gamma)\zeta^{6\nu}z_2z_3+(\gamma-\alpha)\zeta^{2\nu}z_1z_3+\\
&+(\beta-\alpha)\zeta^{8\nu}z_4z_5+(\gamma-\beta)\zeta^{7\nu}z_5z_6+(\alpha-\gamma)\zeta^{11\nu}z_4z_6+\\
&-(\alpha+\beta)\zeta^\nu z_3z_4-(\beta+\gamma)\zeta^{9\nu}z_1z_5-(\gamma+\alpha)\zeta^{3\nu}z_2z_6+\\
&-(\alpha+\beta)\zeta^{12\nu}z_1z_6-(\beta+\gamma)\zeta^{4\nu}z_2z_4-(\gamma+\alpha)\zeta^{10\nu}z_3z_5.
\end{aligned}$$

$$\begin{aligned}
&13ST^\nu(z_2)\cdot ST^\nu(z_5)\\
=&\gamma z_1z_4+\alpha z_2z_5+\beta z_3z_6+\\
&+\alpha\zeta^\nu z_1^2+\beta\zeta^{9\nu}z_2^2+\gamma\zeta^{3\nu}z_3^2-\alpha\zeta^{12\nu}z_4^2-\beta\zeta^{4\nu}z_5^2-\gamma\zeta^{10\nu}z_6^2+\\
&+(\beta-\gamma)\zeta^{5\nu}z_1z_2+(\gamma-\alpha)\zeta^{6\nu}z_2z_3+(\alpha-\beta)\zeta^{2\nu}z_1z_3+\\
&+(\gamma-\beta)\zeta^{8\nu}z_4z_5+(\alpha-\gamma)\zeta^{7\nu}z_5z_6+(\beta-\alpha)\zeta^{11\nu}z_4z_6+\\
&-(\beta+\gamma)\zeta^\nu z_3z_4-(\gamma+\alpha)\zeta^{9\nu}z_1z_5-(\alpha+\beta)\zeta^{3\nu}z_2z_6+\\
&-(\beta+\gamma)\zeta^{12\nu}z_1z_6-(\gamma+\alpha)\zeta^{4\nu}z_2z_4-(\alpha+\beta)\zeta^{10\nu}z_3z_5.
\end{aligned}$$

$$\begin{aligned}
&13ST^\nu(z_3)\cdot ST^\nu(z_6)\\
=&\alpha z_1z_4+\beta z_2z_5+\gamma z_3z_6+\\
&+\beta\zeta^\nu z_1^2+\gamma\zeta^{9\nu}z_2^2+\alpha\zeta^{3\nu}z_3^2-\beta\zeta^{12\nu}z_4^2-\gamma\zeta^{4\nu}z_5^2-\alpha\zeta^{10\nu}z_6^2+\\
&+(\gamma-\alpha)\zeta^{5\nu}z_1z_2+(\alpha-\beta)\zeta^{6\nu}z_2z_3+(\beta-\gamma)\zeta^{2\nu}z_1z_3+\\
&+(\alpha-\gamma)\zeta^{8\nu}z_4z_5+(\beta-\alpha)\zeta^{7\nu}z_5z_6+(\gamma-\beta)\zeta^{11\nu}z_4z_6+\\
&-(\gamma+\alpha)\zeta^\nu z_3z_4-(\alpha+\beta)\zeta^{9\nu}z_1z_5-(\beta+\gamma)\zeta^{3\nu}z_2z_6+\\
&-(\gamma+\alpha)\zeta^{12\nu}z_1z_6-(\alpha+\beta)\zeta^{4\nu}z_2z_4-(\beta+\gamma)\zeta^{10\nu}z_3z_5.
\end{aligned}$$

Note that $\alpha+\beta+\gamma=\sqrt{13}$, we find that

$$\begin{aligned}&\sqrt{13}\left[ST^{\nu}(z_1)\cdot ST^{\nu}(z_4)+ST^{\nu}(z_2)\cdot ST^{\nu}(z_5)+ST^{\nu}(z_3)\cdot ST^{\nu}(z_6)\right]\\=&(z_1z_4+z_2z_5+z_3z_6)+(\zeta^{\nu}z_1^2+\zeta^{9\nu}z_2^2+\zeta^{3\nu}z_3^2)-(\zeta^{12\nu}z_4^2+\zeta^{4\nu}z_5^2+\zeta^{10\nu}z_6^2)+\\&-2(\zeta^{\nu}z_3z_4+\zeta^{9\nu}z_1z_5+\zeta^{3\nu}z_2z_6)-2(\zeta^{12\nu}z_1z_6+\zeta^{4\nu}z_2z_4+\zeta^{10\nu}z_3z_5).\end{aligned}$$

Let

$$\varphi_{\infty}(z_1,z_2,z_3,z_4,z_5,z_6)=\sqrt{13}(z_1z_4+z_2z_5+z_3z_6) \tag{3.1}$$

and

$$\varphi_{\nu}(z_1,z_2,z_3,z_4,z_5,z_6)=\varphi_{\infty}(ST^{\nu}(z_1,z_2,z_3,z_4,z_5,z_6)) \tag{3.2}$$

for $\nu=0,1,\cdots,12$. Then

$$\begin{aligned}\varphi_{\nu}=&(z_1z_4+z_2z_5+z_3z_6)+\zeta^{\nu}(z_1^2-2z_3z_4)+\zeta^{4\nu}(-z_5^2-2z_2z_4)+\\&+\zeta^{9\nu}(z_2^2-2z_1z_5)+\zeta^{3\nu}(z_3^2-2z_2z_6)+\zeta^{12\nu}(-z_4^2-2z_1z_6)+\zeta^{10\nu}(-z_6^2-2z_3z_5).\end{aligned} \tag{3.3}$$

This leads us to define the following senary quadratic forms (quadratic forms in six variables):

$$\begin{cases}\mathbb{A}_0=z_1z_4+z_2z_5+z_3z_6,\\ \mathbb{A}_1=z_1^2-2z_3z_4,\\ \mathbb{A}_2=-z_5^2-2z_2z_4,\\ \mathbb{A}_3=z_2^2-2z_1z_5,\\ \mathbb{A}_4=z_3^2-2z_2z_6,\\ \mathbb{A}_5=-z_4^2-2z_1z_6,\\ \mathbb{A}_6=-z_6^2-2z_3z_5.\end{cases} \tag{3.4}$$

Hence,

$$\sqrt{13}ST^{\nu}(\mathbb{A}_0)=\mathbb{A}_0+\zeta^{\nu}\mathbb{A}_1+\zeta^{4\nu}\mathbb{A}_2+\zeta^{9\nu}\mathbb{A}_3+\zeta^{3\nu}\mathbb{A}_4+\zeta^{12\nu}\mathbb{A}_5+\zeta^{10\nu}\mathbb{A}_6. \tag{3.5}$$

Let

$$\begin{cases}p_1=\zeta^2+\zeta^{11}-2+2(\zeta+\zeta^{12}-\zeta^9-\zeta^4),\\ p_2=2-\zeta^9-\zeta^4+2(\zeta^5+\zeta^8-\zeta^2-\zeta^{11}),\\ p_3=\zeta^6+\zeta^7-2+2(\zeta^3+\zeta^{10}-\zeta-\zeta^{12}),\\ p_4=\zeta^5+\zeta^8-2+2(\zeta^9+\zeta^4-\zeta^3-\zeta^{10}),\\ p_5=2-\zeta^3-\zeta^{10}+2(\zeta^6+\zeta^7-\zeta^5-\zeta^8),\\ p_6=2-\zeta-\zeta^{12}+2(\zeta^2+\zeta^{11}-\zeta^6-\zeta^7).\end{cases} \tag{3.6}$$

We find that

$$\begin{cases} 13S(\mathbb{A}_1) = 2\sqrt{13}\mathbb{A}_0 + p_1\mathbb{A}_1 + p_2\mathbb{A}_2 + p_3\mathbb{A}_3 + p_4\mathbb{A}_4 + p_5\mathbb{A}_5 + p_6\mathbb{A}_6, \\ 13S(\mathbb{A}_2) = 2\sqrt{13}\mathbb{A}_0 + p_2\mathbb{A}_1 + p_4\mathbb{A}_2 + p_6\mathbb{A}_3 + p_5\mathbb{A}_4 + p_3\mathbb{A}_5 + p_1\mathbb{A}_6, \\ 13S(\mathbb{A}_3) = 2\sqrt{13}\mathbb{A}_0 + p_3\mathbb{A}_1 + p_6\mathbb{A}_2 + p_4\mathbb{A}_3 + p_1\mathbb{A}_4 + p_2\mathbb{A}_5 + p_5\mathbb{A}_6, \\ 13S(\mathbb{A}_4) = 2\sqrt{13}\mathbb{A}_0 + p_4\mathbb{A}_1 + p_5\mathbb{A}_2 + p_1\mathbb{A}_3 + p_3\mathbb{A}_4 + p_6\mathbb{A}_5 + p_2\mathbb{A}_6, \\ 13S(\mathbb{A}_5) = 2\sqrt{13}\mathbb{A}_0 + p_5\mathbb{A}_1 + p_3\mathbb{A}_2 + p_2\mathbb{A}_3 + p_6\mathbb{A}_4 + p_1\mathbb{A}_5 + p_4\mathbb{A}_6, \\ 13S(\mathbb{A}_6) = 2\sqrt{13}\mathbb{A}_0 + p_6\mathbb{A}_1 + p_1\mathbb{A}_2 + p_5\mathbb{A}_3 + p_2\mathbb{A}_4 + p_4\mathbb{A}_5 + p_3\mathbb{A}_6, \end{cases} \tag{3.7}$$

Note that

$$\begin{cases} p_1 = \sqrt{13}(\zeta^2 + \zeta^{11}), \\ p_2 = \sqrt{13}(\zeta^9 + \zeta^4), \\ p_3 = \sqrt{13}(\zeta^6 + \zeta^7), \\ p_4 = \sqrt{13}(\zeta^5 + \zeta^8), \\ p_5 = \sqrt{13}(\zeta^3 + \zeta^{10}), \\ p_6 = \sqrt{13}(\zeta + \zeta^{12}). \end{cases} \tag{3.8}$$

We obtain a seven-dimensional representation of the simple group $PSL(2,13) \cong \langle \widetilde{S}, \widetilde{T} \rangle$ which is deduced from the actions of S and T on the basis $(\mathbb{A}_0, \mathbb{A}_1, \mathbb{A}_2, \mathbb{A}_3, \mathbb{A}_4, \mathbb{A}_5, \mathbb{A}_6)$. Here

$$\widetilde{S} = \frac{1}{\sqrt{13}} \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 2 & \zeta^2+\zeta^{11} & \zeta^9+\zeta^4 & \zeta^6+\zeta^7 & \zeta^5+\zeta^8 & \zeta^3+\zeta^{10} & \zeta+\zeta^{12} \\ 2 & \zeta^9+\zeta^4 & \zeta^5+\zeta^8 & \zeta+\zeta^{12} & \zeta^3+\zeta^{10} & \zeta^6+\zeta^7 & \zeta^2+\zeta^{11} \\ 2 & \zeta^6+\zeta^7 & \zeta+\zeta^{12} & \zeta^5+\zeta^8 & \zeta^2+\zeta^{11} & \zeta^9+\zeta^4 & \zeta^3+\zeta^{10} \\ 2 & \zeta^5+\zeta^8 & \zeta^3+\zeta^{10} & \zeta^2+\zeta^{11} & \zeta^6+\zeta^7 & \zeta+\zeta^{12} & \zeta^9+\zeta^4 \\ 2 & \zeta^3+\zeta^{10} & \zeta^6+\zeta^7 & \zeta^9+\zeta^4 & \zeta+\zeta^{12} & \zeta^2+\zeta^{11} & \zeta^5+\zeta^8 \\ 2 & \zeta+\zeta^{12} & \zeta^2+\zeta^{11} & \zeta^3+\zeta^{10} & \zeta^9+\zeta^4 & \zeta^5+\zeta^8 & \zeta^6+\zeta^7 \end{pmatrix}, \tag{3.9}$$

and

$$\widetilde{T} = \begin{pmatrix} 1 & & & & & & \\ & \zeta & & & & & \\ & & \zeta^4 & & & & \\ & & & \zeta^9 & & & \\ & & & & \zeta^3 & & \\ & & & & & \zeta^{12} & \\ & & & & & & \zeta^{10} \end{pmatrix}. \tag{3.10}$$

Now, let us reproduce from [CC] some of the character table of G.

Table 1. Some of the character table of $PSL(2,13)$

	$1A$	$2A$	$3A$	$6A$	$7A$	$7B$	$7C$	$13A$	$13B$
χ_1	1	1	1	1	1	1	1	1	1
χ_2	7	-1	1	-1	0	0	0	$\frac{1-\sqrt{13}}{2}$	$\frac{1+\sqrt{13}}{2}$
χ_3	7	-1	1	-1	0	0	0	$\frac{1+\sqrt{13}}{2}$	$\frac{1-\sqrt{13}}{2}$
χ_{10}	6	0	0	0	-1	-1	-1	$\frac{-1+\sqrt{13}}{2}$	$\frac{-1-\sqrt{13}}{2}$
χ_{11}	6	0	0	0	-1	-1	-1	$\frac{-1-\sqrt{13}}{2}$	$\frac{-1+\sqrt{13}}{2}$
χ_{15}	14	0	2	0	0	0	0	1	1

In [K1], Klein obtained the modular equation of degree fourteen, which corresponds to the transformation of order thirteen:

$$\begin{aligned} J : J-1 : 1 =&(\tau^2+5\tau+13)(\tau^4+7\tau^3+20\tau^2+19\tau+1)^3 \\ &:(\tau^2+6\tau+13)(\tau^6+10\tau^5+46\tau^4+108\tau^3+122\tau^2+38\tau-1)^2 \\ &:1728\tau, \end{aligned}$$

where the Hauptmodul $\tau=(\eta(z)/\eta(13z))^2$. Note that

$$\begin{aligned} &\tau^4+7\tau^3+20\tau^2+19\tau+1 \\ =&\left(\tau^2+\frac{7+\sqrt{13}}{2}\tau+\frac{11+3\sqrt{13}}{2}\right)\left(\tau^2+\frac{7-\sqrt{13}}{2}\tau+\frac{11-3\sqrt{13}}{2}\right), \end{aligned}$$

$$\begin{aligned} &\tau^6+10\tau^5+46\tau^4+108\tau^3+122\tau^2+38\tau-1 \\ =&\left(\tau^3+5\tau^2+\frac{21-\sqrt{13}}{2}\tau+\frac{3+\sqrt{13}}{2}\right)\left(\tau^3+5\tau^2+\frac{21+\sqrt{13}}{2}\tau+\frac{3-\sqrt{13}}{2}\right). \end{aligned}$$

In [De1] and [De2], Deligne gave a modular interpretation of why such factorizations exist. Let us confine our thought to an especially important result which Jacobi had established as early as 1829 in his "Notices sur les fonctions elliptiques" (see [K]). Jacobi there considered, instead of the modular equation, the so-called multiplier-equation, together with other equations equivalent to it, and found that their $(n+1)$ roots are composed in a simple manner of $\frac{n+1}{2}$ elements, with the help of merely numerical irrationalities. Namely, if we denote these elements by $\mathbb{A}_0$, $\mathbb{A}_1$, $\cdots$, $\mathbb{A}_{\frac{n-1}{2}}$, and further, for the roots z of the equation under consideration, apply the indices employed by Galois, we have, with appropriate determination of the square root occurring on the left-hand side:

$$\begin{cases} \sqrt{z_\infty}=\sqrt{(-1)^{\frac{n-1}{2}}\cdot n}\cdot\mathbb{A}_0, \\ \sqrt{z_\nu}=\mathbb{A}_0+\epsilon^\nu\mathbb{A}_1+\epsilon^{4\nu}\mathbb{A}_2+\cdots+\epsilon^{(\frac{n-1}{2})^2\nu}\mathbb{A}_{\frac{n-1}{2}} \end{cases}$$

for $\nu = 0, 1, \cdots, n-1$ and $\epsilon = e^{\frac{2\pi i}{n}}$. Jacobi had himself emphasized the special significance of his result by adding to his short communication: "C'est un théorème des plus importants dans la théorie algébrique de la transformation et de la division des fonctions elliptiques." Now, we give the Jacobian equation of degree fourteen which corresponds to the above modular equation of degree fourteen.

Let $H := Q^5P^2 \cdot P^2Q^6P^8 \cdot Q^5P^2 \cdot P^3Q$ where

$$Q^5P^2 = -\frac{1}{\sqrt{13}} \begin{pmatrix} \zeta^8-\zeta^5 & \zeta^4-\zeta^8 & \zeta^2-\zeta & \zeta^4-\zeta^6 & \zeta^9-\zeta^2 & \zeta-\zeta^6 \\ \zeta^5-\zeta^9 & \zeta^7-\zeta^6 & \zeta^{10}-\zeta^7 & \zeta^9-\zeta^2 & \zeta^{10}-\zeta^2 & \zeta^3-\zeta^5 \\ \zeta^{12}-\zeta^{11} & \zeta^6-\zeta^3 & \zeta^{11}-\zeta^2 & \zeta-\zeta^6 & \zeta^3-\zeta^5 & \zeta^{12}-\zeta^5 \\ \zeta^7-\zeta^9 & \zeta^{11}-\zeta^4 & \zeta^7-\zeta^{12} & \zeta^5-\zeta^8 & \zeta^9-\zeta^5 & \zeta^{11}-\zeta^{12} \\ \zeta^{11}-\zeta^4 & \zeta^{11}-\zeta^3 & \zeta^8-\zeta^{10} & \zeta^8-\zeta^4 & \zeta^6-\zeta^7 & \zeta^3-\zeta^6 \\ \zeta^7-\zeta^{12} & \zeta^8-\zeta^{10} & \zeta^8-\zeta & \zeta-\zeta^2 & \zeta^7-\zeta^{10} & \zeta^2-\zeta^{11} \end{pmatrix},$$

$$P^2Q^6P^8 = -\frac{1}{\sqrt{13}} \begin{pmatrix} \zeta^8-\zeta^5 & \zeta^{12}-\zeta^3 & \zeta^4-\zeta^3 & \zeta^2-\zeta^4 & \zeta^{12}-\zeta^5 & \zeta^{10}-\zeta^2 \\ \zeta^{10}-\zeta & \zeta^7-\zeta^6 & \zeta^4-\zeta & \zeta^{12}-\zeta^5 & \zeta^5-\zeta^{10} & \zeta^4-\zeta^6 \\ \zeta^{10}-\zeta^9 & \zeta^{12}-\zeta^9 & \zeta^{11}-\zeta^2 & \zeta^{10}-\zeta^2 & \zeta^4-\zeta^6 & \zeta^6-\zeta^{12} \\ \zeta^9-\zeta^{11} & \zeta^8-\zeta & \zeta^{11}-\zeta^3 & \zeta^5-\zeta^8 & \zeta-\zeta^{10} & \zeta^9-\zeta^{10} \\ \zeta^8-\zeta & \zeta^3-\zeta^8 & \zeta^7-\zeta^9 & \zeta^3-\zeta^{12} & \zeta^6-\zeta^7 & \zeta^9-\zeta^{12} \\ \zeta^{11}-\zeta^3 & \zeta^7-\zeta^9 & \zeta-\zeta^7 & \zeta^3-\zeta^4 & \zeta-\zeta^4 & \zeta^2-\zeta^{11} \end{pmatrix},$$

and

$$\begin{aligned} &Q^5P^2 \cdot P^2Q^6P^8 \cdot Q^5P^2 \\ &= -\frac{1}{\sqrt{13}} \begin{pmatrix} \zeta^7-\zeta^6 & \zeta^8-\zeta^5 & \zeta^{11}-\zeta^2 & \zeta^4-\zeta^9 & \zeta^{12}-\zeta & \zeta^{10}-\zeta^3 \\ \zeta^8-\zeta^5 & \zeta^{11}-\zeta^2 & \zeta^7-\zeta^6 & \zeta^{12}-\zeta & \zeta^{10}-\zeta^3 & \zeta^4-\zeta^9 \\ \zeta^{11}-\zeta^2 & \zeta^7-\zeta^6 & \zeta^8-\zeta^5 & \zeta^{10}-\zeta^3 & \zeta^4-\zeta^9 & \zeta^{12}-\zeta \\ \zeta^4-\zeta^9 & \zeta^{12}-\zeta & \zeta^{10}-\zeta^3 & \zeta^6-\zeta^7 & \zeta^5-\zeta^8 & \zeta^2-\zeta^{11} \\ \zeta^{12}-\zeta & \zeta^{10}-\zeta^3 & \zeta^4-\zeta^9 & \zeta^5-\zeta^8 & \zeta^2-\zeta^{11} & \zeta^6-\zeta^7 \\ \zeta^{10}-\zeta^3 & \zeta^4-\zeta^9 & \zeta^{12}-\zeta & \zeta^2-\zeta^{11} & \zeta^6-\zeta^7 & \zeta^5-\zeta^8 \end{pmatrix}. \end{aligned}$$

Then

$$H = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix}. \tag{3.11}$$

Note that

$$H^2 = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}, \quad H^3 = \begin{pmatrix} 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

and $H^6=-I$. In the projective coordinates, this means that $H^6=1$. We have

$$H^{-1}TH=-T^4.$$

Thus, $\langle H,T\rangle\cong\mathbb{Z}_{13}\rtimes\mathbb{Z}_6$. Hence, it is a maximal subgroup of order 78 of G with index 14 (see [CC]). We find that φ_∞^2 is invariant under the action of the maximal subgroup $\langle H,T\rangle$. Note that

$$\varphi_\infty=\sqrt{13}\mathbb{A}_0,\quad \varphi_\nu=\mathbb{A}_0+\zeta^\nu\mathbb{A}_1+\zeta^{4\nu}\mathbb{A}_2+\zeta^{9\nu}\mathbb{A}_3+\zeta^{3\nu}\mathbb{A}_4+\zeta^{12\nu}\mathbb{A}_5+\zeta^{10\nu}\mathbb{A}_6$$

for $\nu=0,1,\cdots,12$. Let $w=\varphi^2$, $w_\infty=\varphi_\infty^2$ and $w_\nu=\varphi_\nu^2$. Then w_∞, w_ν for $\nu=0,\cdots,12$ form an algebraic equation of degree fourteen, which is just the Jacobian equation of degree fourteen, whose roots are these w_ν and w_∞:

$$w^{14}+a_1w^{13}+\cdots+a_{13}w+a_{14}=0.$$

In particular, the coefficients

$$a_{14}=\varphi_\infty^2\cdot\prod_{\nu=0}^{12}\varphi_\nu^2=13\mathbb{A}_0^2\prod_{\nu=0}^{12}(\mathbb{A}_0+\zeta^\nu\mathbb{A}_1+\zeta^{4\nu}\mathbb{A}_2+\zeta^{9\nu}\mathbb{A}_3+\zeta^{3\nu}\mathbb{A}_4+\zeta^{12\nu}\mathbb{A}_5+\zeta^{10\nu}\mathbb{A}_6)^2,$$

and

$$-a_1=w_\infty+\sum_{\nu=0}^{12}w_\nu=26(\mathbb{A}_0^2+\mathbb{A}_1\mathbb{A}_5+\mathbb{A}_2\mathbb{A}_3+\mathbb{A}_4\mathbb{A}_6).\tag{3.12}$$

Recall that the modular equation of degree fourteen is also given by (see [Hu])

$$\begin{aligned}&z^{14}+13[2\Delta^{\frac{1}{12}}z^{13}+25\Delta^{\frac{2}{12}}z^{12}+196\Delta^{\frac{3}{12}}z^{11}+1064\Delta^{\frac{4}{12}}z^{10}\\&\quad+4180\Delta^{\frac{5}{12}}z^9+12086\Delta^{\frac{6}{12}}z^8+25660\Delta^{\frac{7}{12}}z^7+39182\Delta^{\frac{8}{12}}z^6\\&\quad+41140\Delta^{\frac{9}{12}}z^5+27272\Delta^{\frac{10}{12}}z^4+9604\Delta^{\frac{11}{12}}z^3+1165\Delta z^2]\\&\quad+[746\Delta-(12g_2)^3]\Delta^{\frac{1}{12}}z+13\Delta^{\frac{14}{12}}=0.\end{aligned}$$

The coefficients of z^{13} and the constant are just 26 and 13.

In fact, the invariant quadric

$$\Psi_2:=\mathbb{A}_0^2+\mathbb{A}_1\mathbb{A}_5+\mathbb{A}_2\mathbb{A}_3+\mathbb{A}_4\mathbb{A}_6=2\Phi_4(z_1,z_2,z_3,z_4,z_5,z_6),\tag{3.13}$$

where

$$\Phi_4:=(z_3z_4^3+z_1z_5^3+z_2z_6^3)-(z_6z_1^3+z_4z_2^3+z_5z_3^3)+3(z_1z_2z_4z_5+z_2z_3z_5z_6+z_3z_1z_6z_4),\tag{3.14}$$

Hence, the variety $\Psi_2 = 0$ is a quartic four-fold, which is invariant under the action of the simple group G.

Recall that the principal congruence subgroup of level 13 is the normal subgroup $\Gamma(13)$ of $\Gamma = PSL(2,\mathbb{Z})$ defined by the exact sequence

$$1 \to \Gamma(13) \to \Gamma(1) \xrightarrow{f} G \to 1 \tag{3.15}$$

where $f(\gamma) \equiv \gamma \pmod{13}$ for $\gamma \in \Gamma = \Gamma(1)$. Then there is a representation

$$\rho : \Gamma \to PGL(6,\mathbb{C}) \tag{3.16}$$

with kernel $\Gamma(13)$ and leaving Φ_4 invariant. It is defined as follows: if $t = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $s = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, then $\rho(t) = T$ and $\rho(s) = S$. To see that such a representation exists, note that Γ is defined by the presentation

$$\langle s, t; s^2 = (st)^3 = 1 \rangle$$

satisfied by s and t and we have proved that S and T satisfy these relations. Moreover, we have proved that G is defined by the presentation

$$\langle S, T; S^2 = T^{13} = (ST)^3 = 1, (Q^3P^4)^3 = 1 \rangle. \tag{3.17}$$

Let $p = st^{-1}s$ and $q = st^3$. Then

$$h := q^5p^2 \cdot p^2q^6p^8 \cdot q^5p^2 \cdot p^3q = \begin{pmatrix} 4,428,249 & -10,547,030 \\ -11,594,791 & 27,616,019 \end{pmatrix} \tag{3.18}$$

satisfies that $\rho(h) = H$.

Theorem 3.1. *The invariant quartic four-fold* $\Phi_4(z_1, z_2, z_3, z_4, z_5, z_6) = 0$ *can be parametrized by the theta constants associated with* $\Gamma(13)$:

$$\Phi_4(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z)) = 0. \tag{3.19}$$

Proof. Let $y_i(z) = \eta^3(z)a_i(z)$ $(1 \le i \le 6)$. Then

$$Y(z) := \begin{pmatrix} y_1(z) \\ y_2(z) \\ y_3(z) \\ y_4(z) \\ y_5(z) \\ y_6(z) \end{pmatrix} = \eta^3(z) \begin{pmatrix} a_1(z) \\ a_2(z) \\ a_3(z) \\ a_4(z) \\ a_5(z) \\ a_6(z) \end{pmatrix} = \eta^3(z)\mathbf{A}(z).$$

It is well-known that $\eta(z)$ satisfies the following transformation formulas

$$\eta(z+1)=e^{\frac{\pi i}{12}}\eta(z),\quad \text{and}\quad \eta\left(-\frac{1}{z}\right)=e^{-\frac{\pi i}{4}}\sqrt{z}\eta(z).$$

By Proposition 2.5, we have

$$Y(z+1)=e^{-\frac{\pi i}{2}}\rho(t)Y(z)\quad \text{and}\quad Y\left(-\frac{1}{z}\right)=e^{-\frac{\pi i}{2}}z^2\rho(s)Y(z).$$

Define $j(\gamma,z):=cz+d$ if $z\in\mathbb{H}$ and $\gamma=\begin{pmatrix}a & b\\ c & d\end{pmatrix}\in\Gamma(1)$. Hence,

$$Y(\gamma(z))=v(\gamma)j(\gamma,z)^2\rho(\gamma)Y(z)\quad \text{for } \gamma\in\Gamma(1),$$

where $v(\gamma)=\pm1$ or $\pm i$. Since $\Gamma(13)=\ker\rho$, we have

$$Y(\gamma(z))=v(\gamma)j(\gamma,z)^2Y(z)\quad \text{for } \gamma\in\Gamma(13).$$

This means that the functions $y_1(z)$, $y_2(z)$, $y_3(z)$, $y_4(z)$, $y_5(z)$, $y_6(z)$ are modular forms of weight two for $\Gamma(13)$ with the same multiplier $v(\gamma)=\pm1$ or $\pm i$. Thus,

$$\Phi_4(Y(\gamma(z)))=v(\gamma)^4j(\gamma,z)^8\Phi_4(Y(z))=j(\gamma,z)^8\Phi_4(Y(z))\quad \text{for } \gamma\in\Gamma(13).$$

Moreover, for $\gamma\in\Gamma(1)$,

$$\begin{aligned}&\Phi_4(Y(\gamma(z)))=\Phi_4(v(\gamma)j(\gamma,z)^2\rho(\gamma)Y(z))\\ =&v(\gamma)^4j(\gamma,z)^8\Phi_4(\rho(\gamma)Y(z))=j(\gamma,z)^8\Phi_4(\rho(\gamma)Y(z)).\end{aligned}$$

Note that $\rho(\gamma)\in\langle\rho(s),\rho(t)\rangle=G$ and Φ_4 is a G-invariant polynomial, we have

$$\Phi_4(Y(\gamma(z)))=j(\gamma,z)^8\Phi_4(Y(z)),\quad \text{for } \gamma\in\Gamma(1).$$

This implies that $\Phi_4(Y(z))$ is a modular form of weight eight for the full modular group $\Gamma(1)$. Moreover, we will show that it is a cusp form. We have

$$a_3(z)a_4(z)^3=-q^{\frac{1}{2}}\sum_{n\in\mathbb{Z}}(-1)^nq^{\frac{1}{2}(13n^2+5n)}\left[\sum_{n\in\mathbb{Z}}(-1)^nq^{\frac{1}{2}(13n^2+3n)}\right]^3,$$

$$a_1(z)a_5(z)^3=q^{\frac{7}{2}}\sum_{n\in\mathbb{Z}}(-1)^nq^{\frac{1}{2}(13n^2+11n)}\left[\sum_{n\in\mathbb{Z}}(-1)^nq^{\frac{1}{2}(13n^2+9n)}\right]^3,$$

$$a_2(z)a_6(z)^3 = q^{\frac{1}{2}} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+7n)} \left[\sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+n)}\right]^3,$$

$$a_6(z)a_1(z)^3 = q^{\frac{7}{2}} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+n)} \left[\sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+11n)}\right]^3,$$

$$a_4(z)a_2(z)^3 = -q^{\frac{3}{2}} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+3n)} \left[\sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+7n)}\right]^3,$$

$$a_5(z)a_3(z)^3 = q^{\frac{3}{2}} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+9n)} \left[\sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+5n)}\right]^3,$$

$$\begin{aligned} a_1(z)a_2(z)a_4(z)a_5(z) =& - q^{\frac{5}{2}} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+11n)} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+7n)} \\ & \times \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+3n)} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+9n)}, \end{aligned}$$

$$\begin{aligned} a_2(z)a_3(z)a_5(z)a_6(z) =& q^{\frac{3}{2}} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+7n)} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+5n)} \\ & \times \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+9n)} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+n)}, \end{aligned}$$

and

$$\begin{aligned} a_3(z)a_1(z)a_6(z)a_4(z) =& - q^{\frac{3}{2}} \sum_{n\in\mathbb{Z}} q^{\frac{1}{2}(13n^2+5n)} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+11n)} \\ & \times \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+n)} \sum_{n\in\mathbb{Z}} (-1)^n q^{\frac{1}{2}(13n^2+3n)}. \end{aligned}$$

Hence,

$$\Phi_4(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z)) = q^{\frac{1}{2}} \sum_{n=0}^{\infty} a_n q^n, \quad \text{where } a_n \in \mathbb{Z}.$$

On the other hand, $\eta(z)^{12} = q^{\frac{1}{2}} \prod_{n=1}^{\infty} (1-q^n)^{12}$. We have

$$\Phi_4(y_1(z), y_2(z), y_3(z), y_4(z), y_5(z), y_6(z)) = q \sum_{n=0}^{\infty} a_n q^n \prod_{n=1}^{\infty} (1-q^n)^{12}$$

is a cusp form of weight 8 for the full modular group $\Gamma = PSL(2,\mathbb{Z})$, but the only such form is zero. (See [Se], Chapter VII for the complete description of cusp forms on $\Gamma(1)$). This completes the proof of Theorem 3.1.

□

Corollary 3.2. *The following invariant decomposition formula* (*exotic modular equation*) *holds:*

$$\frac{\Psi_2(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))}{\mathbb{A}_0(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))^2}=0, \tag{3.20}$$

where the quadric

$$\Psi_2(z_1,z_2,z_3,z_4,z_5,z_6):=\mathbb{A}_0^2+\mathbb{A}_1\mathbb{A}_5+\mathbb{A}_2\mathbb{A}_3+\mathbb{A}_4\mathbb{A}_6 \tag{3.21}$$

is an invariant associated to $PSL(2,13)$ *and* $\mathbb{A}_0^2$ *is invariant under the action of the image of a Borel subgroup of* $PSL(2,13)$ (*i.e. a maximal subgroup of order* 78 *of* $PSL(2,13)$).

Proof. This comes from Theorem 3.1 by noting that (3.13). Thus, we complete the proof of Theorem 1.3.

□

Now, we will prove that

$$\begin{aligned}&\prod_{j=1}^{6}\mathbb{A}_j(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))\\ \neq&-\mathbb{A}_0(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))^6.\end{aligned} \tag{3.22}$$

We have

$$\mathbb{A}_0(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))=q^{\frac{1}{4}}(1+O(q)),$$

$$\mathbb{A}_1(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))=q^{\frac{34}{104}}(2+O(q)),$$

$$\mathbb{A}_2(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))=q^{\frac{58}{104}}(2+O(q)),$$

$$\mathbb{A}_3(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))=q^{\frac{98}{104}}(1+O(q)),$$

$$\mathbb{A}_4(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))=q^{\frac{50}{104}}(-1+O(q)),$$

$$\mathbb{A}_5(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))=q^{\frac{18}{104}}(-1+O(q)),$$

$$\mathbb{A}_6(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))=q^{\frac{2}{104}}(-1+O(q)).$$

Thus,

$$\prod_{j=1}^{6}\mathbb{A}_j(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))=q^{\frac{5}{2}}(-4+O(q)).$$

On the other hand,

$$-\mathbb{A}_0(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))^6 = q^{\frac{3}{2}}(-1+O(q)).$$

This gives the proof of (3.22).

In fact, Theorem 3.1 also gives the geometry of the classical modular curve $X(13)$:

Corollary 3.3. *The coordinates* $(a_1(z),a_2(z),a_3(z),a_4(z),a_5(z),a_6(z))$ *map* $X(13)$ *into the quartic four-fold* $\Phi_4(z_1,z_2,z_3,z_4,z_5,z_6)=0$ *in* $\mathbb{CP}^5$.

Proof. This comes from Theorem 2.3 and Theorem 3.1.

□

4. Fourteen-dimensional representations of $PSL(2,13)$ and invariant decomposition formula

It is known that the principal congruence subgroup $\Gamma(13)$ is a normal subgroup of $\Gamma_0(13)$, and $\Gamma_0(13)$ is generated by $\Gamma(13)$ and the group generated by H and T. The quotient group $\Gamma_0(13)/\Gamma(13)$ is of order 78 and is isomorphic to the subgroup $\langle H,T\rangle$. The natural projection defines a Galois covering $X(13)\to X_0(13)$ with Galois group $\langle H,T\rangle$. Recall that $(\eta(z)/\eta(13z))^2$ is the Hauptmodul of $X_0(13)$, that is, the unique (up to multiplication by a constant) univalent modular function for $\Gamma_0(13)$ holomorphic on $\mathbb{H}$ with a simple pole at $z=i\infty$ and a simple zero at $z=0$.

We will construct a fourteen-dimensional representation of $PSL(2,13)$ which is deduced from our six-dimensional representation. It should be emphasized that our fourteen-dimensional representation is not Weil representation. In contrast with this fact, both our six-dimensional and seven-dimensional representations of G are Weil representations, i.e., $\frac{p-1}{2}$-dimensional and $\frac{p+1}{2}$-dimensional representations of $PSL(2,p)$, respectively. In fact, what Klein used in his papers [K], [K1], [K2], [K3] and [K4] are all Weil representations. Hence, our method is completely different from Klein's method.

To construct our fourteen-dimensional representation which is generated under the action of PSL by a specific vector in Sym^3(six-dimensional representation), we begin

with a cubic polynomial $z_1z_2z_3$ and study the action of ST^ν (ν mod 13) on it. We have

$$
\begin{aligned}
&-13\sqrt{13}ST^\nu(z_1)\cdot ST^\nu(z_2)\cdot ST^\nu(z_3)\\
=&-\sqrt{\frac{-13-3\sqrt{13}}{2}}(\zeta^{8\nu}z_1^3+\zeta^{7\nu}z_2^3+\zeta^{11\nu}z_3^3)-\sqrt{\frac{-13+3\sqrt{13}}{2}}(\zeta^{5\nu}z_4^3+\zeta^{6\nu}z_5^3+\zeta^{2\nu}z_6^3)\\
&-\sqrt{-13+2\sqrt{13}}(\zeta^{12\nu}z_1^2z_2+\zeta^{4\nu}z_2^2z_3+\zeta^{10\nu}z_3^2z_1)\\
&-\sqrt{-13-2\sqrt{13}}(\zeta^{\nu}z_4^2z_5+\zeta^{9\nu}z_5^2z_6+\zeta^{3\nu}z_6^2z_4)\\
&+2\sqrt{-13-2\sqrt{13}}(\zeta^{3\nu}z_1z_2^2+\zeta^{\nu}z_2z_3^2+\zeta^{9\nu}z_3z_1^2)\\
&-2\sqrt{-13+2\sqrt{13}}(\zeta^{10\nu}z_4z_5^2+\zeta^{12\nu}z_5z_6^2+\zeta^{4\nu}z_6z_4^2)\\
&+2\sqrt{\frac{-13-3\sqrt{13}}{2}}(\zeta^{7\nu}z_1^2z_4+\zeta^{11\nu}z_2^2z_5+\zeta^{8\nu}z_3^2z_6)+\\
&-2\sqrt{\frac{-13+3\sqrt{13}}{2}}(\zeta^{6\nu}z_1z_4^2+\zeta^{2\nu}z_2z_5^2+\zeta^{5\nu}z_3z_6^2)+\\
&+\sqrt{-13-2\sqrt{13}}(\zeta^{3\nu}z_1^2z_5+\zeta^{\nu}z_2^2z_6+\zeta^{9\nu}z_3^2z_4)+\\
&+\sqrt{-13+2\sqrt{13}}(\zeta^{10\nu}z_2z_4^2+\zeta^{12\nu}z_3z_5^2+\zeta^{4\nu}z_1z_6^2)+\\
&+\sqrt{\frac{-13+3\sqrt{13}}{2}}(\zeta^{6\nu}z_1^2z_6+\zeta^{2\nu}z_2^2z_4+\zeta^{5\nu}z_3^2z_5)+\\
&+\sqrt{\frac{-13-3\sqrt{13}}{2}}(\zeta^{7\nu}z_3z_4^2+\zeta^{11\nu}z_1z_5^2+\zeta^{8\nu}z_2z_6^2)+\\
&+[2(\theta_1-\theta_3)-3(\theta_2-\theta_4)]z_1z_2z_3+[2(\theta_4-\theta_2)-3(\theta_1-\theta_3)]z_4z_5z_6+
\end{aligned}
$$

$$
\begin{aligned}
&-\sqrt{\frac{-13-3\sqrt{13}}{2}}(\zeta^{11\nu}z_1z_2z_4+\zeta^{8\nu}z_2z_3z_5+\zeta^{7\nu}z_1z_3z_6)+\\
&+\sqrt{\frac{-13+3\sqrt{13}}{2}}(\zeta^{2\nu}z_1z_4z_5+\zeta^{5\nu}z_2z_5z_6+\zeta^{6\nu}z_3z_4z_6)+\\
&-3\sqrt{\frac{-13-3\sqrt{13}}{2}}(\zeta^{7\nu}z_1z_2z_5+\zeta^{11\nu}z_2z_3z_6+\zeta^{8\nu}z_1z_3z_4)+\\
&+3\sqrt{\frac{-13+3\sqrt{13}}{2}}(\zeta^{6\nu}z_2z_4z_5+\zeta^{2\nu}z_3z_5z_6+\zeta^{5\nu}z_1z_4z_6)+\\
&-\sqrt{-13+2\sqrt{13}}(\zeta^{10\nu}z_1z_2z_6+\zeta^{4\nu}z_1z_3z_5+\zeta^{12\nu}z_2z_3z_4)+\\
&+\sqrt{-13-2\sqrt{13}}(\zeta^{3\nu}z_3z_4z_5+\zeta^{9\nu}z_2z_4z_6+\zeta^{\nu}z_1z_5z_6).
\end{aligned}
$$

This leads us to define the following senary cubic forms (cubic forms in six variables):

$$
\begin{cases}
\mathbb{D}_0=z_1z_2z_3,\\
\mathbb{D}_1=2z_2z_3^2+z_2^2z_6-z_4^2z_5+z_1z_5z_6,\\
\mathbb{D}_2=-z_6^3+z_2^2z_4-2z_2z_5^2+z_1z_4z_5+3z_3z_5z_6,\\
\mathbb{D}_3=2z_1z_2^2+z_1^2z_5-z_4z_6^2+z_3z_4z_5,\\
\mathbb{D}_4=-z_2^2z_3+z_1z_6^2-2z_4^2z_6-z_1z_3z_5,\\
\mathbb{D}_5=-z_4^3+z_3^2z_5-2z_3z_6^2+z_2z_5z_6+3z_1z_4z_6,\\
\mathbb{D}_6=-z_5^3+z_1^2z_6-2z_1z_4^2+z_3z_4z_6+3z_2z_4z_5,\\
\mathbb{D}_7=-z_2^3+z_3z_4^2-z_1z_3z_6-3z_1z_2z_5+2z_1^2z_4,\\
\mathbb{D}_8=-z_1^3+z_2z_6^2-z_2z_3z_5-3z_1z_3z_4+2z_3^2z_6,\\
\mathbb{D}_9=2z_1^2z_3+z_3^2z_4-z_5^2z_6+z_2z_4z_6,\\
\mathbb{D}_{10}=-z_1z_3^2+z_2z_4^2-2z_4z_5^2-z_1z_2z_6,\\
\mathbb{D}_{11}=-z_3^3+z_1z_5^2-z_1z_2z_4-3z_2z_3z_6+2z_2^2z_5,\\
\mathbb{D}_{12}=-z_1^2z_2+z_3z_5^2-2z_5z_6^2-z_2z_3z_4,\\
\mathbb{D}_{\infty}=z_4z_5z_6.
\end{cases}
\tag{4.1}
$$

Let

$$r_0=2(\theta_1-\theta_3)-3(\theta_2-\theta_4),\quad r_\infty=2(\theta_4-\theta_2)-3(\theta_1-\theta_3),$$

and

$$r_1=\sqrt{-13-2\sqrt{13}},r_2=\sqrt{\frac{-13+3\sqrt{13}}{2}},r_3=\sqrt{-13+2\sqrt{13}},r_4=\sqrt{\frac{-13-3\sqrt{13}}{2}}.$$

Then

$$\begin{aligned}&-13\sqrt{13}ST^{\nu}(\mathbb{D}_0)\\ =&r_0\mathbb{D}_0+r_1\zeta^{\nu}\mathbb{D}_1+r_2\zeta^{2\nu}\mathbb{D}_2+r_1\zeta^{3\nu}\mathbb{D}_3+r_3\zeta^{4\nu}\mathbb{D}_4+r_2\zeta^{5\nu}\mathbb{D}_5+r_2\zeta^{6\nu}\mathbb{D}_6+\\ &+r_4\zeta^{7\nu}\mathbb{D}_7+r_4\zeta^{8\nu}\mathbb{D}_8+r_1\zeta^{9\nu}\mathbb{D}_9+r_3\zeta^{10\nu}\mathbb{D}_{10}+r_4\zeta^{11\nu}\mathbb{D}_{11}+r_3\zeta^{12\nu}\mathbb{D}_{12}+r_\infty\mathbb{D}_\infty.\end{aligned}$$

$$\begin{aligned}-13\sqrt{13}S(\mathbb{D}_\infty)=&r_\infty\mathbb{D}_0-r_3\mathbb{D}_1-r_4\mathbb{D}_2-r_3\mathbb{D}_3+r_1\mathbb{D}_4-r_4\mathbb{D}_5-r_4\mathbb{D}_6+\\ &+r_2\mathbb{D}_7+r_2\mathbb{D}_8-r_3\mathbb{D}_9+r_1\mathbb{D}_{10}+r_2\mathbb{D}_{11}+r_1\mathbb{D}_{12}-r_0\mathbb{D}_\infty.\end{aligned}$$

Moreover, we have

$$\begin{aligned}-13\sqrt{13}S(\mathbb{D}_1)=&13r_1\mathbb{D}_0+q_1\mathbb{D}_1+q_2\mathbb{D}_2+q_3\mathbb{D}_3+q_4\mathbb{D}_4+q_5\mathbb{D}_5+q_6\mathbb{D}_6+\\ &+q_7\mathbb{D}_7+q_8\mathbb{D}_8+q_9\mathbb{D}_9+q_{10}\mathbb{D}_{10}+q_{11}\mathbb{D}_{11}+q_{12}\mathbb{D}_{12}-13r_3\mathbb{D}_\infty,\end{aligned}$$

where

$$\begin{cases}q_1=-2(\zeta-\zeta^{12})-2(\zeta^5-\zeta^8)+6(\zeta^3-\zeta^{10})-(\zeta^2-\zeta^{11})+4(\zeta^9-\zeta^4)+2(\zeta^6-\zeta^7),\\ q_2=-4(\zeta-\zeta^{12})+3(\zeta^5-\zeta^8)+3(\zeta^3-\zeta^{10})-(\zeta^2-\zeta^{11})-2(\zeta^9-\zeta^4),\\ q_3=6(\zeta-\zeta^{12})-(\zeta^5-\zeta^8)+4(\zeta^3-\zeta^{10})+2(\zeta^2-\zeta^{11})-2(\zeta^9-\zeta^4)-2(\zeta^6-\zeta^7),\\ q_4=-2(\zeta-\zeta^{12})+4(\zeta^5-\zeta^8)+2(\zeta^3-\zeta^{10})-2(\zeta^2-\zeta^{11})+(\zeta^9-\zeta^4)+6(\zeta^6-\zeta^7),\\ q_5=-2(\zeta-\zeta^{12})-4(\zeta^3-\zeta^{10})+3(\zeta^2-\zeta^{11})+3(\zeta^9-\zeta^4)-(\zeta^6-\zeta^7),\\ q_6=3(\zeta-\zeta^{12})-(\zeta^5-\zeta^8)-2(\zeta^3-\zeta^{10})-4(\zeta^9-\zeta^4)+3(\zeta^6-\zeta^7),\\ q_7=(\zeta-\zeta^{12})+3(\zeta^5-\zeta^8)-2(\zeta^2-\zeta^{11})-3(\zeta^9-\zeta^4)-4(\zeta^6-\zeta^7),\\ q_8=-2(\zeta^5-\zeta^8)-3(\zeta^3-\zeta^{10})-4(\zeta^2-\zeta^{11})+(\zeta^9-\zeta^4)+3(\zeta^6-\zeta^7),\\ q_9=4(\zeta-\zeta^{12})+2(\zeta^5-\zeta^8)-2(\zeta^3-\zeta^{10})-2(\zeta^2-\zeta^{11})+6(\zeta^9-\zeta^4)-(\zeta^6-\zeta^7),\\ q_{10}=(\zeta-\zeta^{12})+6(\zeta^5-\zeta^8)-2(\zeta^3-\zeta^{10})+4(\zeta^2-\zeta^{11})+2(\zeta^9-\zeta^4)-2(\zeta^6-\zeta^7),\\ q_{11}=-3(\zeta-\zeta^{12})-4(\zeta^5-\zeta^8)+(\zeta^3-\zeta^{10})+3(\zeta^2-\zeta^{11})-2(\zeta^6-\zeta^7),\\ q_{12}=2(\zeta-\zeta^{12})-2(\zeta^5-\zeta^8)+(\zeta^3-\zeta^{10})+6(\zeta^2-\zeta^{11})-2(\zeta^9-\zeta^4)+4(\zeta^6-\zeta^7).\end{cases}\tag{4.2}$$

Similarly, we obtain

$$\begin{aligned}-13\sqrt{13}S(\mathbb{D}_2)=&26r_2\mathbb{D}_0+2q_2\mathbb{D}_1-q_4\mathbb{D}_2+2q_6\mathbb{D}_3+2q_8\mathbb{D}_4-q_{10}\mathbb{D}_5-q_{12}\mathbb{D}_6+\\ &+q_1\mathbb{D}_7+q_3\mathbb{D}_8+2q_5\mathbb{D}_9+2q_7\mathbb{D}_{10}+q_9\mathbb{D}_{11}+2q_{11}\mathbb{D}_{12}-26r_4\mathbb{D}_\infty.\end{aligned}$$

$$\begin{aligned}-13\sqrt{13}S(\mathbb{D}_3)=&13r_1\mathbb{D}_0+q_3\mathbb{D}_1+q_6\mathbb{D}_2+q_9\mathbb{D}_3+q_{12}\mathbb{D}_4+q_2\mathbb{D}_5+q_5\mathbb{D}_6+\\ &+q_8\mathbb{D}_7+q_{11}\mathbb{D}_8+q_1\mathbb{D}_9+q_4\mathbb{D}_{10}+q_7\mathbb{D}_{11}+q_{10}\mathbb{D}_{12}-13r_3\mathbb{D}_\infty,\end{aligned}$$

$$
\begin{aligned}
-13\sqrt{13}S(\mathbb{D}_4) =& 13r_3\mathbb{D}_0 + q_4\mathbb{D}_1 + q_8\mathbb{D}_2 + q_{12}\mathbb{D}_3 - q_3\mathbb{D}_4 + q_7\mathbb{D}_5 + q_{11}\mathbb{D}_6 + \\
& - q_2\mathbb{D}_7 - q_6\mathbb{D}_8 + q_{10}\mathbb{D}_9 - q_1\mathbb{D}_{10} - q_5\mathbb{D}_{11} - q_9\mathbb{D}_{12} + 13r_1\mathbb{D}_\infty, \\
-13\sqrt{13}S(\mathbb{D}_5) =& 26r_2\mathbb{D}_0 + 2q_5\mathbb{D}_1 - q_{10}\mathbb{D}_2 + 2q_2\mathbb{D}_3 + 2q_7\mathbb{D}_4 - q_{12}\mathbb{D}_5 - q_4\mathbb{D}_6 + \\
& + q_9\mathbb{D}_7 + q_1\mathbb{D}_8 + 2q_6\mathbb{D}_9 + 2q_{11}\mathbb{D}_{10} + q_3\mathbb{D}_{11} + 2q_8\mathbb{D}_{12} - 26r_4\mathbb{D}_\infty. \\
-13\sqrt{13}S(\mathbb{D}_6) =& 26r_2\mathbb{D}_0 + 2q_6\mathbb{D}_1 - q_{12}\mathbb{D}_2 + 2q_5\mathbb{D}_3 + 2q_{11}\mathbb{D}_4 - q_4\mathbb{D}_5 - q_{10}\mathbb{D}_6 + \\
& + q_3\mathbb{D}_7 + q_9\mathbb{D}_8 + 2q_2\mathbb{D}_9 + 2q_8\mathbb{D}_{10} + q_1\mathbb{D}_{11} + 2q_7\mathbb{D}_{12} - 26r_4\mathbb{D}_\infty. \\
-13\sqrt{13}S(\mathbb{D}_7) =& 26r_4\mathbb{D}_0 + 2q_7\mathbb{D}_1 + q_1\mathbb{D}_2 + 2q_8\mathbb{D}_3 - 2q_2\mathbb{D}_4 + q_9\mathbb{D}_5 + q_3\mathbb{D}_6 + \\
& + q_{10}\mathbb{D}_7 + q_4\mathbb{D}_8 + 2q_{11}\mathbb{D}_9 - 2q_5\mathbb{D}_{10} + q_{12}\mathbb{D}_{11} - 2q_6\mathbb{D}_{12} + 26r_2\mathbb{D}_\infty. \\
-13\sqrt{13}S(\mathbb{D}_8) =& 26r_4\mathbb{D}_0 + 2q_8\mathbb{D}_1 + q_3\mathbb{D}_2 + 2q_{11}\mathbb{D}_3 - 2q_6\mathbb{D}_4 + q_1\mathbb{D}_5 + q_9\mathbb{D}_6 + \\
& + q_4\mathbb{D}_7 + q_{12}\mathbb{D}_8 + 2q_7\mathbb{D}_9 - 2q_2\mathbb{D}_{10} + q_{10}\mathbb{D}_{11} - 2q_5\mathbb{D}_{12} + 26r_2\mathbb{D}_\infty. \\
-13\sqrt{13}S(\mathbb{D}_9) =& 13r_1\mathbb{D}_0 + q_9\mathbb{D}_1 + q_5\mathbb{D}_2 + q_1\mathbb{D}_3 + q_{10}\mathbb{D}_4 + q_6\mathbb{D}_5 + q_2\mathbb{D}_6 + \\
& + q_1\mathbb{D}_7 + q_7\mathbb{D}_8 + q_8\mathbb{D}_9 + q_{12}\mathbb{D}_{10} + q_8\mathbb{D}_{11} + q_4\mathbb{D}_{12} - 13r_3\mathbb{D}_\infty, \\
-13\sqrt{13}S(\mathbb{D}_{10}) =& 13r_3\mathbb{D}_0 + q_{10}\mathbb{D}_1 + q_7\mathbb{D}_2 + q_4\mathbb{D}_3 - q_1\mathbb{D}_4 + q_{11}\mathbb{D}_5 + q_8\mathbb{D}_6 + \\
& - q_5\mathbb{D}_7 - q_2\mathbb{D}_8 + q_{12}\mathbb{D}_9 - q_9\mathbb{D}_{10} - q_6\mathbb{D}_{11} - q_3\mathbb{D}_{12} + 13r_1\mathbb{D}_\infty, \\
-13\sqrt{13}S(\mathbb{D}_{11}) =& 26r_4\mathbb{D}_0 + 2q_{11}\mathbb{D}_1 + q_9\mathbb{D}_2 + 2q_7\mathbb{D}_3 - 2q_5\mathbb{D}_4 + q_3\mathbb{D}_5 + q_1\mathbb{D}_6 + \\
& + q_{12}\mathbb{D}_7 + q_{10}\mathbb{D}_8 + 2q_8\mathbb{D}_9 - 2q_6\mathbb{D}_{10} + q_4\mathbb{D}_{11} - 2q_2\mathbb{D}_{12} + 26r_2\mathbb{D}_\infty. \\
-13\sqrt{13}S(\mathbb{D}_{12}) =& 13r_3\mathbb{D}_0 + q_{12}\mathbb{D}_1 + q_{11}\mathbb{D}_2 + q_{10}\mathbb{D}_3 - q_9\mathbb{D}_4 + q_8\mathbb{D}_5 + q_7\mathbb{D}_6 + \\
& - q_6\mathbb{D}_7 - q_5\mathbb{D}_8 + q_4\mathbb{D}_9 - q_3\mathbb{D}_{10} - q_2\mathbb{D}_{11} - q_1\mathbb{D}_{12} + 13r_1\mathbb{D}_\infty.
\end{aligned}
$$

Hence, we get the element $\widehat{S}$ which is induced from the action of S on the basis $(\mathbb{D}_0, \cdots, \mathbb{D}_\infty)$:

$$
\widehat{S} = -\frac{1}{13\sqrt{13}} \begin{pmatrix} S_1 & S_2 \\ S_3 & S_4 \end{pmatrix}, \tag{4.3}
$$

with

$$
S_1 = \begin{pmatrix}
r_0 & r_1 & r_2 & r_1 & r_3 & r_2 & r_2 \\
13r_1 & q_1 & q_2 & q_3 & q_4 & q_5 & q_6 \\
26r_2 & 2q_2 & -q_4 & 2q_6 & 2q_8 & -q_{10} & -q_{12} \\
13r_1 & q_3 & q_6 & q_9 & q_{12} & q_2 & q_5 \\
13r_3 & q_4 & q_8 & q_{12} & -q_3 & q_7 & q_{11} \\
26r_2 & 2q_5 & -q_{10} & 2q_2 & 2q_7 & -q_{12} & -q_4 \\
26r_2 & 2q_6 & -q_{12} & 2q_5 & 2q_{11} & -q_4 & -q_{10}
\end{pmatrix}, \tag{4.4}
$$

$$S_2 = \begin{pmatrix} r_4 & r_4 & r_1 & r_3 & r_4 & r_3 & r_\infty \\ q_7 & q_8 & q_9 & q_{10} & q_{11} & q_{12} & -13r_3 \\ q_1 & q_3 & 2q_5 & 2q_7 & q_9 & 2q_{11} & -26r_4 \\ q_8 & q_{11} & q_1 & q_4 & q_7 & q_{10} & -13r_3 \\ -q_2 & -q_6 & q_{10} & -q_1 & -q_5 & -q_9 & 13r_1 \\ q_9 & q_1 & 2q_6 & 2q_{11} & q_3 & 2q_8 & -26r_4 \\ q_3 & q_9 & 2q_2 & 2q_8 & q_1 & 2q_7 & -26r_4 \end{pmatrix}, \tag{4.5}$$

$$S_3 = \begin{pmatrix} 26r_4 & 2q_7 & q_1 & 2q_8 & -2q_2 & q_9 & q_3 \\ 26r_4 & 2q_8 & q_3 & 2q_{11} & -2q_6 & q_1 & q_9 \\ 13r_1 & q_9 & q_5 & q_1 & q_{10} & q_6 & q_2 \\ 13r_3 & q_{10} & q_7 & q_4 & -q_1 & q_{11} & q_8 \\ 26r_4 & 2q_{11} & q_9 & 2q_7 & -2q_5 & q_3 & q_1 \\ 13r_3 & q_{12} & q_{11} & q_{10} & -q_9 & q_8 & q_7 \\ r_\infty & -r_3 & -r_4 & -r_3 & r_1 & -r_4 & -r_4 \end{pmatrix}, \tag{4.6}$$

$$S_4 = \begin{pmatrix} q_{10} & q_4 & 2q_{11} & -2q_5 & q_{12} & -2q_6 & 26r_2 \\ q_4 & q_{12} & 2q_7 & -2q_2 & q_{10} & -2q_5 & 26r_2 \\ q_{11} & q_7 & q_3 & q_{12} & q_8 & q_4 & -13r_3 \\ -q_5 & -q_2 & q_{12} & -q_9 & -q_6 & -q_3 & 13r_1 \\ q_{12} & q_{10} & 2q_8 & -2q_6 & q_4 & -2q_2 & 26r_2 \\ -q_6 & -q_5 & q_4 & -q_3 & -q_2 & -q_1 & 13r_1 \\ r_2 & r_2 & -r_3 & r_1 & r_2 & r_1 & -r_0 \end{pmatrix}. \tag{4.7}$$

Similarly, the element $\widehat{T}$ is induced from the action of T on the basis $(\mathbb{D}_0, \cdots, \mathbb{D}_\infty)$:

$$\widehat{T} = \mathrm{diag}(1, \zeta, \zeta^2, \zeta^3, \zeta^4, \zeta^5, \zeta^6, \zeta^7, \zeta^8, \zeta^9, \zeta^{10}, \zeta^{11}, \zeta^{12}, 1). \tag{4.8}$$

We have

$$\mathrm{Tr}(\widehat{S}) = 0, \quad \mathrm{Tr}(\widehat{T}) = 1, \quad \mathrm{Tr}(\widehat{S}\widehat{T}) = -2. \tag{4.9}$$

Hence, this fourteen-dimensional representation corresponds to the character χ_{15} of $PSL(2, 13)$ in [CC]. It is defined over the cyclotomic field $\mathbb{Q}(\zeta)$.

Let

$$\delta_\infty(z_1, z_2, z_3, z_4, z_5, z_6) = 13^2(z_1^2 z_2^2 z_3^2 + z_4^2 z_5^2 z_6^2) \tag{4.10}$$

and

$$\delta_\nu(z_1, z_2, z_3, z_4, z_5, z_6) = \delta_\infty(ST^\nu(z_1, z_2, z_3, z_4, z_5, z_6)) \tag{4.11}$$

for $\nu = 0, 1, \cdots, 12$. Then

$$\delta_\nu = 13^2 ST^\nu(\mathbb{G}_0) = -13\mathbb{G}_0 + \zeta^\nu \mathbb{G}_1 + \zeta^{2\nu}\mathbb{G}_2 + \cdots + \zeta^{12\nu}\mathbb{G}_{12}, \tag{4.12}$$

where the senary sextic forms (i.e., sextic forms in six variables) are given as follows:

$$\begin{cases}
\mathbb{G}_0 = \mathbb{D}_0^2 + \mathbb{D}_\infty^2, \\
\mathbb{G}_1 = -\mathbb{D}_7^2 + 2\mathbb{D}_0\mathbb{D}_1 + 10\mathbb{D}_\infty\mathbb{D}_1 + 2\mathbb{D}_2\mathbb{D}_{12} - 2\mathbb{D}_3\mathbb{D}_{11} - 4\mathbb{D}_4\mathbb{D}_{10} - 2\mathbb{D}_9\mathbb{D}_5, \\
\mathbb{G}_2 = -2\mathbb{D}_1^2 - 4\mathbb{D}_0\mathbb{D}_2 + 6\mathbb{D}_\infty\mathbb{D}_2 - 2\mathbb{D}_4\mathbb{D}_{11} + 2\mathbb{D}_5\mathbb{D}_{10} - 2\mathbb{D}_6\mathbb{D}_9 - 2\mathbb{D}_7\mathbb{D}_8, \\
\mathbb{G}_3 = -\mathbb{D}_8^2 + 2\mathbb{D}_0\mathbb{D}_3 + 10\mathbb{D}_\infty\mathbb{D}_3 + 2\mathbb{D}_6\mathbb{D}_{10} - 2\mathbb{D}_9\mathbb{D}_7 - 4\mathbb{D}_{12}\mathbb{D}_4 - 2\mathbb{D}_1\mathbb{D}_2, \\
\mathbb{G}_4 = -\mathbb{D}_2^2 + 10\mathbb{D}_0\mathbb{D}_4 - 2\mathbb{D}_\infty\mathbb{D}_4 + 2\mathbb{D}_5\mathbb{D}_{12} - 2\mathbb{D}_9\mathbb{D}_8 - 4\mathbb{D}_1\mathbb{D}_3 - 2\mathbb{D}_{10}\mathbb{D}_7, \\
\mathbb{G}_5 = -2\mathbb{D}_9^2 - 4\mathbb{D}_0\mathbb{D}_5 + 6\mathbb{D}_\infty\mathbb{D}_5 - 2\mathbb{D}_{10}\mathbb{D}_8 + 2\mathbb{D}_6\mathbb{D}_{12} - 2\mathbb{D}_2\mathbb{D}_3 - 2\mathbb{D}_{11}\mathbb{D}_7, \\
\mathbb{G}_6 = -2\mathbb{D}_3^2 - 4\mathbb{D}_0\mathbb{D}_6 + 6\mathbb{D}_\infty\mathbb{D}_6 - 2\mathbb{D}_{12}\mathbb{D}_7 + 2\mathbb{D}_2\mathbb{D}_4 - 2\mathbb{D}_5\mathbb{D}_1 - 2\mathbb{D}_8\mathbb{D}_{11}, \\
\mathbb{G}_7 = -2\mathbb{D}_{10}^2 + 6\mathbb{D}_0\mathbb{D}_7 + 4\mathbb{D}_\infty\mathbb{D}_7 - 2\mathbb{D}_1\mathbb{D}_6 - 2\mathbb{D}_2\mathbb{D}_5 - 2\mathbb{D}_8\mathbb{D}_{12} - 2\mathbb{D}_9\mathbb{D}_{11}, \\
\mathbb{G}_8 = -2\mathbb{D}_4^2 + 6\mathbb{D}_0\mathbb{D}_8 + 4\mathbb{D}_\infty\mathbb{D}_8 - 2\mathbb{D}_3\mathbb{D}_5 - 2\mathbb{D}_6\mathbb{D}_2 - 2\mathbb{D}_{11}\mathbb{D}_{10} - 2\mathbb{D}_1\mathbb{D}_7, \\
\mathbb{G}_9 = -\mathbb{D}_{11}^2 + 2\mathbb{D}_0\mathbb{D}_9 + 10\mathbb{D}_\infty\mathbb{D}_9 + 2\mathbb{D}_5\mathbb{D}_4 - 2\mathbb{D}_1\mathbb{D}_8 - 4\mathbb{D}_{10}\mathbb{D}_{12} - 2\mathbb{D}_3\mathbb{D}_6, \\
\mathbb{G}_{10} = -\mathbb{D}_5^2 + 10\mathbb{D}_0\mathbb{D}_{10} - 2\mathbb{D}_\infty\mathbb{D}_{10} + 2\mathbb{D}_6\mathbb{D}_4 - 2\mathbb{D}_3\mathbb{D}_7 - 4\mathbb{D}_9\mathbb{D}_1 - 2\mathbb{D}_{12}\mathbb{D}_{11}, \\
\mathbb{G}_{11} = -2\mathbb{D}_{12}^2 + 6\mathbb{D}_0\mathbb{D}_{11} + 4\mathbb{D}_\infty\mathbb{D}_{11} - 2\mathbb{D}_9\mathbb{D}_2 - 2\mathbb{D}_5\mathbb{D}_6 - 2\mathbb{D}_7\mathbb{D}_4 - 2\mathbb{D}_3\mathbb{D}_8, \\
\mathbb{G}_{12} = -\mathbb{D}_6^2 + 10\mathbb{D}_0\mathbb{D}_{12} - 2\mathbb{D}_\infty\mathbb{D}_{12} + 2\mathbb{D}_2\mathbb{D}_{10} - 2\mathbb{D}_1\mathbb{D}_{11} - 4\mathbb{D}_3\mathbb{D}_9 - 2\mathbb{D}_4\mathbb{D}_8.
\end{cases} \tag{4.13}$$

We have that $\mathbb{G}_0$ is invariant under the action of $\langle H, T\rangle$, a maximal subgroup of order 78 of G with index 14. Note that δ_∞, δ_ν for $\nu = 0, \cdots, 12$ form an algebraic equation of degree fourteen. However, we have $\delta_\infty + \sum_{\nu=0}^{12} \delta_\nu = 0$. Hence, it is not the Jacobian equation of degree fourteen! We call it exotic modular equation of degree fourteen. We have

$$\delta_\infty^2 + \sum_{\nu=0}^{12} \delta_\nu^2 = 26(7 \cdot 13^2 \mathbb{G}_0^2 + \mathbb{G}_1\mathbb{G}_{12} + \mathbb{G}_2\mathbb{G}_{11} + \cdots + \mathbb{G}_6\mathbb{G}_7). \tag{4.14}$$

This leads us to define

$$\Phi_{12}(z_1, z_2, z_3, z_4, z_5, z_6) := -\frac{1}{26}(7 \cdot 13^2 \mathbb{G}_0^2 + \mathbb{G}_1\mathbb{G}_{12} + \mathbb{G}_2\mathbb{G}_{11} + \cdots + \mathbb{G}_6\mathbb{G}_7), \tag{4.15}$$

Theorem 4.1. *The invariant decomposition formula for the simple group $PSL(2, 13)$ of order* 1092 *is given as follows:*

$$\left[\frac{\eta^2(z)}{\eta^2(13z)}\right]^5 = \frac{\Phi_{12}(a_1(z), a_2(z), a_3(z), a_4(z), a_5(z), a_6(z))}{(a_1(z)a_2(z)a_3(z)a_4(z)a_5(z)a_6(z))^2}, \tag{4.16}$$

where $\Phi_{12}(z_1, z_2, z_3, z_4, z_5, z_6)$ is an invariant of degree 12 *associated to $PSL(2, 13)$, and $(z_1z_2z_3z_4z_5z_6)^2$ is invariant under the action of the image of a Borel subgroup of $PSL(2, 13)$ (i.e. a maximal subgroup of order* 78 *of $PSL(2, 13)$).*

Proof. Let $x_i(z) = \eta(z)a_i(z)$ $(1 \leq i \leq 6)$. Then

$$X(z) := \begin{pmatrix} x_1(z) \\ x_2(z) \\ x_3(z) \\ x_4(z) \\ x_5(z) \\ x_6(z) \end{pmatrix} = \eta(z) \begin{pmatrix} a_1(z) \\ a_2(z) \\ a_3(z) \\ a_4(z) \\ a_5(z) \\ a_6(z) \end{pmatrix} = \eta(z)\mathbf{A}(z).$$

Recall that $\eta(z)$ satisfies the following transformation formulas

$$\eta(z+1) = e^{\frac{\pi i}{12}}\eta(z), \quad \text{and} \quad \eta\left(-\frac{1}{z}\right) = e^{-\frac{\pi i}{4}}\sqrt{z}\eta(z).$$

By Proposition 2.5, we have

$$X(z+1) = e^{-\frac{2\pi i}{3}}\rho(t)X(z) \quad \text{and} \quad X\left(-\frac{1}{z}\right) = z\rho(s)X(z).$$

Hence,

$$X(\gamma(z)) = u(\gamma)j(\gamma, z)\rho(\gamma)X(z) \quad \text{for } \gamma \in \Gamma(1),$$

where $u(\gamma) = 1, \omega$ or ω^2 with $\omega = e^{\frac{2\pi i}{3}}$. Since $\Gamma(13) = \ker \rho$, we have

$$X(\gamma(z)) = u(\gamma)j(\gamma, z)X(z) \quad \text{for } \gamma \in \Gamma(13).$$

This means that the functions $x_1(z)$, $x_2(z)$, $x_3(z)$, $x_4(z)$, $x_5(z)$, $x_6(z)$ are modular forms of weight one for $\Gamma(13)$ with the same multiplier $u(\gamma) = 1, \omega$ or ω^2. Thus,

$$\Phi_{12}(X(\gamma(z))) = u(\gamma)^{12}j(\gamma, z)^{12}\Phi_{12}(X(z)) = j(\gamma, z)^{12}\Phi_{12}(X(z)) \quad \text{for } \gamma \in \Gamma(13).$$

Moreover, for $\gamma \in \Gamma(1)$,

$$\begin{aligned} &\Phi_{12}(X(\gamma(z))) = \Phi_{12}(u(\gamma)j(\gamma, z)\rho(\gamma)X(z)) \\ =&u(\gamma)^{12}j(\gamma, z)^{12}\Phi_{12}(\rho(\gamma)X(z)) = j(\gamma, z)^{12}\Phi_{12}(\rho(\gamma)X(z)). \end{aligned}$$

Note that $\rho(\gamma) \in \langle \rho(s), \rho(t) \rangle = G$ and Φ_{12} is a G-invariant polynomial, we have

$$\Phi_{12}(X(\gamma(z))) = j(\gamma, z)^{12}\Phi_{12}(X(z)), \quad \gamma \in \Gamma(1).$$

This implies that $\Phi_{12}(X(z))$ is a modular form of weight 12 for the full modular group $\Gamma(1)$. Moreover, we will show that it is a cusp form. We have

$$\begin{cases} \mathbb{D}_0 = q^{\frac{15}{8}}(1+O(q)), \\ \mathbb{D}_\infty = q^{\frac{7}{8}}(-1+O(q)), \\ \mathbb{D}_1 = q^{\frac{99}{104}}(2+O(q)), \\ \mathbb{D}_2 = q^{\frac{3}{104}}(-1+O(q)), \\ \mathbb{D}_3 = q^{\frac{11}{104}}(1+O(q)), \\ \mathbb{D}_4 = q^{\frac{19}{104}}(-2+O(q)), \\ \mathbb{D}_5 = q^{\frac{27}{104}}(-1+O(q)), \end{cases} \qquad \begin{cases} \mathbb{D}_6 = q^{\frac{35}{104}}(-1+O(q)), \\ \mathbb{D}_7 = q^{\frac{43}{104}}(1+O(q)), \\ \mathbb{D}_8 = q^{\frac{51}{104}}(3+O(q)), \\ \mathbb{D}_9 = q^{\frac{59}{104}}(-2+O(q)), \\ \mathbb{D}_{10} = q^{\frac{67}{104}}(1+O(q)), \\ \mathbb{D}_{11} = q^{\frac{75}{104}}(-4+O(q)), \\ \mathbb{D}_{12} = q^{\frac{83}{104}}(-1+O(q)). \end{cases}$$

Hence,

$$\begin{cases} \mathbb{G}_0 = q^{\frac{7}{4}}(1+O(q)), \\ \mathbb{G}_1 = q^{\frac{86}{104}}(13+O(q)), \\ \mathbb{G}_2 = q^{\frac{94}{104}}(-22+O(q)), \\ \mathbb{G}_3 = q^{\frac{102}{104}}(-21+O(q)), \\ \mathbb{G}_4 = q^{\frac{6}{104}}(-1+O(q)), \\ \mathbb{G}_5 = q^{\frac{14}{104}}(2+O(q)), \\ \mathbb{G}_6 = q^{\frac{22}{104}}(2+O(q)), \end{cases} \qquad \begin{cases} \mathbb{G}_7 = q^{\frac{30}{104}}(-2+O(q)), \\ \mathbb{G}_8 = q^{\frac{38}{104}}(-8+O(q)), \\ \mathbb{G}_9 = q^{\frac{46}{104}}(6+O(q)), \\ \mathbb{G}_{10} = q^{\frac{54}{104}}(1+O(q)), \\ \mathbb{G}_{11} = q^{\frac{62}{104}}(-8+O(q)), \\ \mathbb{G}_{12} = q^{\frac{70}{104}}(17+O(q)). \end{cases}$$

Therefore,

$$\begin{aligned} &7\cdot 13^2\mathbb{G}_0^2 + \mathbb{G}_1\mathbb{G}_{12} + \mathbb{G}_2\mathbb{G}_{11} + \cdots + \mathbb{G}_6\mathbb{G}_7 \\ =&7\cdot 13^2 q^{\frac{7}{2}}(1+O(q)) + q^{\frac{3}{2}}(13\cdot 17+O(q)) + q^{\frac{3}{2}}(22\cdot 8+O(q)) + q^{\frac{3}{2}}(-21+O(q)) + \\ &+ q^{\frac{1}{2}}(-6+O(q)) + q^{\frac{1}{2}}(-16+O(q)) + q^{\frac{1}{2}}(-4+O(q)) \\ =&q^{\frac{1}{2}}(-26+O(q)). \end{aligned}$$

We have

$$\begin{aligned} &\Phi_{12}(x_1(z), x_2(z), x_3(z), x_4(z), x_5(z), x_6(z)) \\ =&\eta(z)^{12} q^{\frac{1}{2}}(1+O(q)) \\ =&q^{\frac{1}{2}}\prod_{n=1}^{\infty}(1-q^n)^{12}\cdot q^{\frac{1}{2}}(1+O(q)) \\ =&q(1+O(q)) \end{aligned}$$

is a cusp form of weight 12 for the full modular group $\Gamma(1)$. Because every $\Gamma(1)$ cusp form of weight 12 is a multiple of $\Delta(z)$, checking the q^1 coefficient, we find that $\Phi_{12}(x_1(z), x_2(z), x_3(z), x_4(z), x_5(z), x_6(z)) = \Delta(z)$. On the other hand, we have

$$x_1(z)x_2(z)x_3(z)x_4(z)x_5(z)x_6(z) = -\eta(z)^7\eta(13z)^5.$$

This completes the proof of Theorem 4.1. □

Note that (2.20), (2.21) and (2.22) are equivalent to the following:

$$\begin{aligned} &a_1(z)^2a_4(z)^2a_2(z)a_5(z) + a_2(z)^2a_5(z)^2a_3(z)a_6(z) + a_3(z)^2a_6(z)^2a_1(z)a_4(z)+ \\ &- a_1(z)a_4(z)a_2(z)a_5(z)a_3(z)a_6(z) = -\eta(z)^3\eta(13z)^3, \end{aligned} \tag{4.17}$$

$$\begin{aligned} &a_1(z)^2a_4(z)^2a_3(z)a_6(z) + a_2(z)^2a_5(z)^2a_1(z)a_4(z) + a_3(z)^2a_6(z)^2a_2(z)a_5(z)+ \\ &+ 4a_1(z)a_4(z)a_2(z)a_5(z)a_3(z)a_6(z) = \eta(z)^3\eta(13z)^3, \end{aligned} \tag{4.18}$$

$$\begin{aligned} &a_4(z)^2a_5(z)^2a_6(z)^2 - a_1(z)^2a_2(z)^2a_3(z)^2 + 3a_1(z)a_4(z)a_2(z)a_5(z)a_3(z)a_6(z) \\ =&\eta(z)^3\eta(13z)^3. \end{aligned} \tag{4.19}$$

The corresponding polynomials are given by

$$f_6(z_1, z_2, z_3, z_4, z_5, z_6) := z_1^2z_4^2z_2z_5 + z_2^2z_5^2z_3z_6 + z_3^2z_6^2z_1z_4 - z_1z_2z_3z_4z_5z_6,$$

$$g_6(z_1, z_2, z_3, z_4, z_5, z_6) := z_1^2z_4^2z_3z_6 + z_2^2z_5^2z_1z_4 + z_3^2z_6^2z_2z_5 + 4z_1z_2z_3z_4z_5z_6,$$

$$h_6(z_1, z_2, z_3, z_4, z_5, z_6) := z_4^2z_5^2z_6^2 - z_1^2z_2^2z_3^2 + 3z_1z_2z_3z_4z_5z_6.$$

Similar as the above argument in the proof of Theorem 4.1, we have

$$\begin{aligned} &f_6(X(\gamma(z))) = f_6(u(\gamma)j(\gamma, z)\rho(\gamma)X(z)) \\ =&u(\gamma)^6j(\gamma, z)^6f_6(\rho(\gamma)X(z)) = j(\gamma, z)^6f_6(\rho(\gamma)X(z)) \quad \text{for } \gamma \in \Gamma(1). \end{aligned}$$

If f_6 is a G-invariant polynomial, we have

$$f_6(X(\gamma(z))) = j(\gamma, z)^6f_6(X(z)) \quad \text{for } \gamma \in \Gamma(1).$$

This implies that $f_6(X(z))$ is a modular form of weight 6 for the full modular group $\Gamma(1)$. On the other hand, by (4.17), we have

$$f_6(X(z)) = -\eta(z)^9\eta(13z)^3 = -q^2\prod_{n=1}^{\infty}(1-q^n)^9\prod_{n=1}^{\infty}(1-q^{13n})^3.$$

This shows that $f_6(X(z))$ is a cusp form of weight 6 for the full modular group $\Gamma(1)$, but the only such form is zero. This leads to a contradiction! Therefore, f_6 is not a G-invariant polynomial. Similarly, we can prove that g_6 and h_6 are not G-invariant polynomials.

Appendix

In this appendix, we give some importance of (1.1) and (1.2) from another point of view. According to [G2], we have the following tables:

Table 1. η-products in $S_{\frac{1}{2}(p-1)}(p, \chi_p)$ for small p

p	η-product cusp forms
5	–
7	$\eta(z)^3\eta(7z)^3$
11	–
13	$\eta(z)^{2n+1}\eta(13z)^{11-2n} \quad (0 \le n \le 5)$

Table 2. Dimension of certain spaces of cusp forms

p	$\dim S_{\frac{1}{2}(p-1)}(p, \chi_p)$
5	0
7	1
11	3
13	6

Note that

$$\begin{aligned} q^{\frac{19}{24}}\sum_{n=0}^{\infty} p(5n+4)q^n &= \frac{1}{\eta(z)^5}\cdot 0 + 5\frac{\eta(5z)^5}{\eta(z)^6}, \\ q^{\frac{17}{24}}\sum_{n=0}^{\infty} p(7n+5)q^n &= \frac{1}{\eta(z)^7}\cdot 7\eta(z)^3\eta(7z)^3 + 49\frac{\eta(7z)^7}{\eta(z)^8}, \end{aligned} \tag{A.1}$$

$$\begin{aligned} q^{\frac{11}{24}}\sum_{n=0}^{\infty} p(13n+6)q^n =& \frac{1}{\eta(z)^{13}}[11\eta(13z)\eta(z)^{11} + 36\cdot 13\eta(13z)^3\eta(z)^9 + \\ &+ 38\cdot 13^2\eta(13z)^5\eta(z)^7 + 20\cdot 13^3\eta(13z)^7\eta(z)^5 + \\ &+ 6\cdot 13^4\eta(13z)^9\eta(z)^3 + 13^5\eta(13z)^{11}\eta(z)] + 13^5\frac{\eta(13z)^{13}}{\eta(z)^{14}}. \end{aligned} \tag{A.2}$$

We observe that what appear in the right-hand side of (A.1) and (A.2) are just the base vectors of $S_{\frac{1}{2}(p-1)}(p, \chi_p)$ for $p = 5$, 7 and 13.

In the end, let us recall some basic fact about the Monster finite simple group $\mathbb{M}$ (see [CN])and its relation with its type $5B$, $7B$ and $13B$ elements. For each prime p with $(p-1)|24$ there is a conjugacy class of elements of $\mathbb{M}$, with centraliser of form $p^{1+2d}.G_p$, where $p.G_p$ is the centraliser of a corresponding automorphism of the Leech Lattice. The symbol p^{1+2d} denotes an extraspecial p-group, and $2d = 24/(p-1)$. Some maximal p-local subgroups of $\mathbb{M}$ are given as follows (see [CC]):

p	structure	specification
2	$2_+^{1+24} \cdot Co_1$	$N(2B)$
3	$3_+^{1+12} \cdot 2Suz:2$	$N(3B)$
5	$5_+^{1+6} : 4J_2 \cdot 2$	$N(5B)$
7	$7_+^{1+4} : (3 \times 2S_7)$	$N(7B)$
13	$13_+^{1+2} : (3 \times 4S_4)$	$N(13B)$

and

p	structure	specification
2	$2^2.2^{11}.2^{22}.(S_3 \times M_{24})$	$N(2B^2)$
3	$3^2.3^5.3^{10}.(M_{11} \times 2S_4)$	$N(3B^2)$
5	$5^2.5^2.5^4.(S_3 \times GL_2(5))$	$N(5B^2)$
7	$7^2.7.7^2 : GL_2(7)$	$N(7B^2)$
13	$13^2 : 4L_2(13) \cdot 2$	$N(13B^2)$

Department of Mathematics, Peking University
Beijing 100871, P. R. China
E-mail address: yanglei@math.pku.edu.cn

References

[AB] S. Ahlgren and M. Boylan, Arithmetic properties of the partition function, Invent. Math. **153** (2003), 487-502.

[AG] G. E. Andrews and F. G. Garvan, Dyson's crank of a partition, Bull. Amer. Math. Soc. **18** (1988), 167-171.

[AB1] G. E. Andrews and B. C. Berndt, *Ramanujan's Lost Notebook*, Part I, Springer, 2005.

[ASD] A. O. L. Atkin and H. P. F. Swinnerton-Dyer, Some properties of partitions, Proc. London Math. Soc. **66** (1954), 84-106.

[B1] B. C. Berndt, *Ramanujan's Notebooks*, Part III, Springer-Verlag, 1991.

[B2] B. C. Berndt, *Ramanujan's Notebooks*, Part IV, Springer-Verlag, 1994.

[BCCL] B. C. Berndt, H. H. Chan, S. H. Chan and W.-C. Liaw, Cranks–really, the final problem, Ramanujan J. **23** (2010), 3-15.

[BO] B. C. Berndt and K. Ono, Ramanujan's unpublished manuscript on the partition and tau functions with proofs and commentary, in: *The Andrews Festschrift, Seventeen Papers on Classical Number Theory and Combinatorics*, 39-110, Springer, 2001.

[Bor] R. E. Borcherds, Monstrous moonshine and monstrous Lie superalgebras, Invent. Math. **109** (1992), 405-444.

[BrO] K. Bringmann and K. Ono, Dyson's ranks and Maass forms, Ann. of Math. (2) **171** (2010), 419-449.

[CY] I. Chen and N. Yui, Singular values of Thompson series, In: *Groups, difference sets, and the Monster (Columbus, OH, 1993)*, 255-326, Ohio State Univ. Math. Res. Inst. Publ., 4, de Gruyter, Berlin, 1996.

[CN] J. Conway and S. Norton, Monstrous moonshine, Bull. London Math. Soc. **11** (1979), 308-339.

[CC] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of Finite Groups, Maximal Subgroups and Ordinary Characters for Simple Groups*, Clarendon Press, Oxford, 1985.

[De1] P. Deligne, Letter to the author, June 29, 2014. Private communication.

[De2] P. Deligne, Letter to the author, August 7, 2014. Private communication.

[Du] W. Duke, Continued fractions and modular functions, Bull. Amer. Math. Soc. (N.S.) **42** (2005), 137-162.

[D] F. J. Dyson, Some guesses in the theory of partitions, Eureka (Cambridge) **8** (1944), 10-15.

[El] N. D. Elkies, The Klein quartic in number theory, *The eightfold way*, 51-101, Math. Sci. Res. Inst. Publ., **35**, Cambridge Univ. Press, Cambridge, 1999.

[Ev] R. J. Evans, Theta function identities, J. Math. Anal. Appl. **147** (1990), 97-121.

[FK] H. M. Farkas and I. Kra, *Theta Constants, Riemann Surfaces and the Modular Group, An Introduction with Applications to Uniformization Theorems, Partition Identities and Combinatorial Number Theory*, Graduate Studies in Mathematics, **37**, American Mathematical Society, Providence, RI, 2001.

[G1] F. G. Garvan, New combinatorial interpretations of Ramanujan's partition congruences mod 5, 7 and 11, Trans. Amer. Math. Soc. **305** (1988), 47-77.

[G2] F. G. Garvan, Some congruences for partitions that are p-cores, Proc. London Math. Soc. **66** (1993), 449-478.

[GKS] F. Garvan, D. Kim and D. Stanton, Cranks and t-cores, Invent. Math. **101** (1990), 1-17.

[Gr] B. H. Gross, Heegner points and the modular curve of prime level, J. Math. Soc. Japan **39** (1987), 345-362.

[Hu] A. Hurwitz, Grundlagen einer independenten Theorie der elliptischen Modulfunktionen und Theorie der Multiplikator-Gleichungen erster Stufe, Math. Ann. **18** (1881), 528-592, in: *Mathematische Werke*, Bd. 1, 1-66, E. Birkhäuser, 1932-33.

[K] F. Klein, *Lectures on the Icosahedron and the Solution of Equations of the Fifth Degree*, Translated by G. G. Morrice, second and revised edition, Dover Publications, Inc., 1956.

[K1] F. Klein, Ueber die Transformation der elliptischen Functionen und die Auflösung der Gleichungen fünften Grades, Math. Ann. **14** (1879), 111-172, in: *Gesammelte Mathematische Abhandlungen*, Bd. III, 13-75, Springer-Verlag, Berlin, 1923.

[K2] F. Klein, Ueber die Transformation siebenter Ordnung der elliptischen Functionen, Math. Ann. **14** (1879), 428-471, in: *Gesammelte Mathematische Abhandlungen*, Bd. III, 90-136, Springer-Verlag, Berlin, 1923.

[K3] F. Klein, Ueber die Auflösung gewisser Gleichungen vom siebenten und achten Grade, Math. Ann. **15** (1879), 251-282, in: *Gesammelte Mathematische Abhandlungen*, Bd. II, 390-438, Springer-Verlag, Berlin, 1922.

[K4] F. Klein, Ueber die Transformation elfter Ordnung der elliptischen Functionen, Math. Ann. **15** (1879), 533-555, in: *Gesammelte Mathematische Abhandlungen*, Bd. III, 140-168, Springer-Verlag, Berlin, 1923.

[KF1] F. Klein and R. Fricke, *Vorlesungen über die Theorie der Elliptischen Modulfunctionen*, Vol. I, Leipzig, 1890.

[KF2] F. Klein and R. Fricke, *Vorlesungen über die Theorie der Elliptischen Modulfunctionen*, Vol. II, Leipzig, 1892.

[KL] D. S. Kubert and S. Lang, *Modular Units*, Grundlehren der Mathematischen Wissenschaften **244**, Springer-Verlag, 1981.

[La] G. Lachaud, Ramanujan modular forms and the Klein quartic, Mosc. Math. J. **5** (2005), 829-856.

[Ma] K. Mahlburg, Partition congruences and the Andrews-Garvan-Dyson crank, Proc. Natl. Acad. Sci. USA **102** (2005), 15373-15376.

[O] K. Ono, Distribution of the partition function modulo m, Ann. of Math. (2) **151** (2000), 293-307.

[Ra] H. Rademacher, The Ramanujan identities under modular substitutions, Trans. Amer. Math. Soc. **51** (1942), 609-636.

[R1] S. Ramanujan, Some properties of $p(n)$, the number of partitions of n, Proc. Cambridge Phil. Soc. **19** (1919), 207-210, in: *Collected Papers of Srinivasa Ramanujan*, 210-213, Cambridge, 1927.

[R2] S. Ramanujan, Congruence properties of partitions, Proc. London Math. Soc. **18** (1920), 19, in: *Collected Papers of Srinivasa Ramanujan*, 230, Cambridge, 1927.

[R3] S. Ramanujan, Congruence properties of partitions, Math. Z. **9** (1921), 147-153, in: *Collected Papers of Srinivasa Ramanujan*, 232-238, Cambridge, 1927.

[R4] S. Ramanujan, Algebraic relations between certain infinite products, Proc. London Math. Soc. **18** (1920), in: *Collected Papers of Srinivasa Ramanujan*, 231, Cambridge, 1927.

[R5] S. Ramanujan, *Notebooks of Srinivasa Ramanujan*, Vol. I, Springer-Verlag, 1984.

[R6] S. Ramanujan, *Notebooks of Srinivasa Ramanujan*, Vol. II, Springer-Verlag, 1984.

[R] S. Ramanujan, *The Lost Notebook and Other Unpublished Papers*, Narosa, New Delhi, 1988.

[Se] J.-P. Serre, *A Course in Arithmetic*, Graduate Texts in Math. **7**, Springer, New York, 1973.

[S] A. Sinkov, Necessary and sufficient conditions for generating certain simple groups by two operators of periods two and three, Amer. J. Math. **59** (1937), 67-76.

[W] A. Weil, Sur certains groupes d'opérateurs unitaires, Acta Math. **111** (1964), 143-211.

[Y] L. Yang, Exotic arithmetic structure on the first Hurwitz triplet. arXiv:1209.1783v5 [math.NT], 2013.

[Z] H. S. Zuckerman, Identities analogous to Ramanujan's identities involving the partition function, Duke Math. J. **5** (1939), 88-110.